

ADAMTS5 Antibody (C-Terminus)
Rabbit Polyclonal Antibody
Catalog # ALS10910**Specification**

ADAMTS5 Antibody (C-Terminus) - Product Information

Application	IHC-P
Primary Accession	Q9UNA0
Reactivity	Human, Monkey
Host	Rabbit
Clonality	Polyclonal
Calculated MW	102kDa KDa
Dilution	IHC-P~~N/A

ADAMTS5 Antibody (C-Terminus) - Additional Information**Gene ID** 11096**Other Names**

A disintegrin and metalloproteinase with thrombospondin motifs 5, ADAM-TS 5, ADAM-TS5, ADAMTS-5, 3.4.24.-, A disintegrin and metalloproteinase with thrombospondin motifs 11, ADAM-TS 11, ADAMTS-11, ADMP-2, AggreCANase-2, ADAMTS5, ADAMTS11, ADMP2

Target/Specificity

Human ADAMTS5. BLAST analysis of the peptide immunogen showed no homology with other human proteins.

Reconstitution & Storage

Long term: -70°C; Short term: +4°C

Precautions

ADAMTS5 Antibody (C-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

ADAMTS5 Antibody (C-Terminus) - Protein Information**Name** ADAMTS5**Synonyms** ADAMTS11, ADMP2**Function**

Metalloproteinase that plays an important role in connective tissue organization, development, inflammation and cell migration. Extracellular matrix (ECM) degrading enzyme that show proteolytic activity toward the hyaluronan group of chondroitin sulfate proteoglycans (CSPGs) including ACAN, VCAN, BCAN and NCAN (PubMed:16133547, PubMed:18992360). Cleavage within the hyaluronans occurs at Glu-Xaa recognition motifs. Plays a role in embryonic development,

including limb and cardiac morphogenesis, and skeletal muscle development through its VCAN remodeling properties. Cleaves VCAN in the pericellular matrix surrounding myoblasts, facilitating myoblast contact and fusion which is required for skeletal muscle development and regeneration (By similarity). Participates in development of brown adipose tissue and browning of white adipose tissue (By similarity). Plays an important role for T-lymphocyte migration from draining lymph nodes following viral infection.

Cellular Location

Secreted, extracellular space, extracellular matrix

Tissue Location

Expressed at low level in placenta primarily but also detected in heart and brain, cervix, uterus, bladder, esophagus, rib cartilage, chondroblastoma, fibrous tissue and a joint capsule from an arthritic patient


Volume

50 µl


ADAMTS5 Antibody (C-Terminus) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ADAMTS5 Antibody (C-Terminus) - Images

Anti-ADAMTS5 antibody ALS10910 IHC of human cartilage.



Anti-ADAMTS5 antibody IHC of human Lung, Non-Small Cell Carcinoma.

ADAMTS5 Antibody (C-Terminus) - Background

Cleaves aggrecan, a cartilage proteoglycan, and may be involved in its turnover. May play an important role in the destruction of aggrecan in arthritic diseases. May play a role in proteolytic processing mostly during the peri-implantation period.

ADAMTS5 Antibody (C-Terminus) - References

- Abbaszade I.,et al.J. Biol. Chem. 274:23443-23450(1999).
Hattori M.,et al.Nature 405:311-319(2000).
Hurskainen T.L.,et al.J. Biol. Chem. 274:25555-25563(1999).
Wang L.W.,et al.J. Biol. Chem. 284:30004-30015(2009).
Mosyak L.,et al.Protein Sci. 17:16-21(2008).