

ERP44 Antibody (Internal)
Rabbit Polyclonal Antibody
Catalog # ALS10959**Specification**

ERP44 Antibody (Internal) - Product Information

Application	IHC
Primary Accession	Q9BS26
Reactivity	Human, Monkey
Host	Rabbit
Clonality	Polyclonal
Calculated MW	47kDa KDa

ERP44 Antibody (Internal) - Additional Information**Gene ID** 23071**Other Names**

Endoplasmic reticulum resident protein 44, ER protein 44, ERp44, Thioredoxin domain-containing protein 4, ERP44, KIAA0573, TXNDC4

Target/Specificity

Human ERP44 / TXNDC4. BLAST analysis of the peptide immunogen showed no homology with other human proteins.

Reconstitution & Storage

Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles.

Precautions

ERP44 Antibody (Internal) is for research use only and not for use in diagnostic or therapeutic procedures.

ERP44 Antibody (Internal) - Protein Information**Name** ERP44**Synonyms** KIAA0573, TXNDC4**Function**

Mediates thiol-dependent retention in the early secretory pathway, forming mixed disulfides with substrate proteins through its conserved CRFS motif (PubMed: 11847130, PubMed: 14517240). Inhibits the calcium channel activity of ITPR1 (PubMed: 15652484). May have a role in the control of oxidative protein folding in the endoplasmic reticulum (PubMed: 11847130, PubMed: 29858230, PubMed: 14517240).

target="_blank">14517240). Required to retain ERO1A and ERO1B in the endoplasmic reticulum (PubMed:11847130, PubMed:29858230).

Cellular Location

Endoplasmic reticulum lumen


Volume

50 µl

ERP44 Antibody (Internal) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ERP44 Antibody (Internal) - Images

Anti-ERP44 / TXNDC4 antibody ALS10959 IHC of human stomach.

ERP44 Antibody (Internal) - Background

Mediates thiol-dependent retention in the early secretory pathway, forming mixed disulfides with substrate proteins through its conserved CRFS motif. Inhibits the calcium channel activity of ITPR1. May have a role in the control of oxidative protein folding in the endoplasmic reticulum. Required to retain ERO1L and ERO1LB in the endoplasmic reticulum.

ERP44 Antibody (Internal) - References

Anelli T., et al. EMBO J. 21:835-844(2002).
Nagase T., et al. DNA Res. 5:31-39(1998).
Clark H.F., et al. Genome Res. 13:2265-2270(2003).
Ota T., et al. Nat. Genet. 36:40-45(2004).

Humphray S.J.,et al.Nature 429:369-374(2004).