

**RICTOR Antibody (C-Terminus)**  
**Goat Polyclonal Antibody**  
**Catalog # ALS11206****Specification**

---

**RICTOR Antibody (C-Terminus) - Product Information**

Application	IHC
Primary Accession	<a href="#">Q6R327</a>
Reactivity	Human, Monkey, Horse, Dog
Host	Goat
Clonality	Polyclonal
Calculated MW	192kDa KDa

**RICTOR Antibody (C-Terminus) - Additional Information****Gene ID** 253260**Other Names**

Rapamycin-insensitive companion of mTOR, AVO3 homolog, hAVO3, RICTOR  
{ECO:0000312|EMBL:EAW55980.1}

**Target/Specificity**

Human RICTOR.

**Reconstitution & Storage**

Store at -20°C. Minimize freezing and thawing.

**Precautions**

RICTOR Antibody (C-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

**RICTOR Antibody (C-Terminus) - Protein Information****Name** RICTOR ([HGNC:28611](#))**Function**


Subunit of mTORC2, which regulates cell growth and survival in response to hormonal signals. mTORC2 is activated by growth factors, but, in contrast to mTORC1, seems to be nutrient-insensitive. mTORC2 seems to function upstream of Rho GTPases to regulate the actin cytoskeleton, probably by activating one or more Rho-type guanine nucleotide exchange factors. mTORC2 promotes the serum-induced formation of stress-fibers or F-actin. mTORC2 plays a critical role in AKT1 'Ser-473' phosphorylation, which may facilitate the phosphorylation of the activation loop of AKT1 on 'Thr-308' by PDK1 which is a prerequisite for full activation. mTORC2 regulates the phosphorylation of SGK1 at 'Ser-422'. mTORC2 also modulates the phosphorylation of PRKCA on 'Ser-657'. Plays an essential role in embryonic growth and development.

## **RICTOR Antibody (C-Terminus) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## **RICTOR Antibody (C-Terminus) - Images**



Anti-RICTOR antibody IHC of human renal tubular epithelium.

## **RICTOR Antibody (C-Terminus) - Background**

Subunit of mTORC2, which regulates cell growth and survival in response to hormonal signals. mTORC2 is activated by growth factors, but, in contrast to mTORC1, seems to be nutrient-insensitive. mTORC2 seems to function upstream of Rho GTPases to regulate the actin cytoskeleton, probably by activating one or more Rho-type guanine nucleotide exchange factors. mTORC2 promotes the serum-induced formation of stress-fibers or F-actin. mTORC2 plays a critical role in AKT1 'Ser-473' phosphorylation, which may facilitate the phosphorylation of the activation loop of AKT1 on 'Thr-308' by PDK1 which is a prerequisite for full activation. mTORC2 regulates the phosphorylation of SGK1 at 'Ser-422'. mTORC2 also modulates the phosphorylation of PRKCA on 'Ser-657'. Plays an essential role in embryonic growth and development.

## **RICTOR Antibody (C-Terminus) - References**

Sarbassov D.D., et al. Curr. Biol. 14:1296-1302(2004).  
Bechtel S., et al. BMC Genomics 8:399-399(2007).  
Schmutz J., et al. Nature 431:268-274(2004).  
Mural R.J., et al. Submitted (JUL-2005) to the EMBL/GenBank/DBJ databases.  
Ohara O., et al. DNA Res. 9:47-57(2002).