

AGAP2 / PIKE Antibody (C-Terminus)
Rabbit Polyclonal Antibody
Catalog # ALS11263**Specification**

AGAP2 / PIKE Antibody (C-Terminus) - Product Information

Application	IHC
Primary Accession	Q99490
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	125kDa KDa

AGAP2 / PIKE Antibody (C-Terminus) - Additional Information**Gene ID** 116986**Other Names**

Arf-GAP with GTPase, ANK repeat and PH domain-containing protein 2, AGAP-2, Centaurin-gamma-1, Cnt-g1, GTP-binding and GTPase-activating protein 2, GGAP2, Phosphatidylinositol 3-kinase enhancer, PIKE, AGAP2, CENTG1, KIAA0167

Target/Specificity

Anti-PIKE antibody will detect both PIKE-L and PIKE-A isoforms.

Reconstitution & Storage

Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles. Store undiluted.

Precautions

AGAP2 / PIKE Antibody (C-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

AGAP2 / PIKE Antibody (C-Terminus) - Protein Information**Name** AGAP2**Synonyms** CENTG1, KIAA0167**Function**

GTPase-activating protein (GAP) for ARF1 and ARF5, which also shows strong GTPase activity. Isoform 1 participates in the prevention of neuronal apoptosis by enhancing PI3 kinase activity. It aids the coupling of metabotropic glutamate receptor 1 (GRM1) to cytoplasmic PI3 kinase by interacting with Homer scaffolding proteins, and also seems to mediate anti-apoptotic effects of NGF by activating nuclear PI3 kinase. Isoform 2 does not stimulate PI3 kinase but may protect cells from apoptosis by stimulating Akt. It also regulates the adapter protein 1 (AP-1)-dependent trafficking of proteins in the endosomal system. It seems to be oncogenic. It is overexpressed in cancer cells, prevents apoptosis and promotes cancer cell invasion.

Cellular Location

[Isoform 1]: Cytoplasm. Nucleus.


Tissue Location

Isoform 1 is brain-specific. Isoform 2 is ubiquitously expressed, with highest levels in brain and heart

AGAP2 / PIKE Antibody (C-Terminus) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

AGAP2 / PIKE Antibody (C-Terminus) - Images

Anti-AGAP2 / PIKE antibody IHC of human heart.

AGAP2 / PIKE Antibody (C-Terminus) - Background

GTPase-activating protein (GAP) for ARF1 and ARF5, which also shows strong GTPase activity. Isoform 1 participates in the prevention of neuronal apoptosis by enhancing PI3 kinase activity. It aids the coupling of metabotropic glutamate receptor 1 (GRM1) to cytoplasmic PI3 kinase by interacting with Homer scaffolding proteins, and also seems to mediate anti-apoptotic effects of NGF by activating nuclear PI3 kinase. Isoform 2 does not stimulate PI3 kinase but may protect cells from apoptosis by stimulating Akt. It also regulates the adapter protein 1 (AP-1)-dependent trafficking of proteins in the endosomal system. It seems to be oncogenic. It is overexpressed in cancer cells, prevents apoptosis and promotes cancer cell invasion.

AGAP2 / PIKE Antibody (C-Terminus) - References

Elkahloun A.G., et al. Genomics 42:295-301(1997).
Roe B., et al. Submitted (JAN-2002) to the EMBL/GenBank/DDBJ databases.
Xia C., et al. Mol. Cell. Biol. 23:2476-2488(2003).

Rong R.,et al.Nat. Neurosci. 6:1153-1161(2003).

Hong W.,et al.Submitted (AUG-2001) to the EMBL/GenBank/DDBJ databases.