

VDAC1 / PORIN Antibody (aa185-197)
Rabbit Polyclonal Antibody
Catalog # ALS11368

Specification

VDAC1 / PORIN Antibody (aa185-197) - Product Information

Application	IHC
Primary Accession	P21796
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	31kDa KDa

VDAC1 / PORIN Antibody (aa185-197) - Additional Information

Gene ID 7416

Other Names

Voltage-dependent anion-selective channel protein 1, VDAC-1, hVDAC1, Outer mitochondrial membrane protein porin 1, Plasmalemmal porin, Porin 31HL, Porin 31HM, VDAC1, VDAC

Target/Specificity

Amino acids 185-197 of Human VDAC1/Porin1.

Reconstitution & Storage

+4°C or -20°C, Avoid repeated freezing and thawing.

Precautions

VDAC1 / PORIN Antibody (aa185-197) is for research use only and not for use in diagnostic or therapeutic procedures.

VDAC1 / PORIN Antibody (aa185-197) - Protein Information

Name VDAC1

Synonyms VDAC

Function

Forms a channel through the mitochondrial outer membrane and also the plasma membrane. The channel at the outer mitochondrial membrane allows diffusion of small hydrophilic molecules; in the plasma membrane it is involved in cell volume regulation and apoptosis. It adopts an open conformation at low or zero membrane potential and a closed conformation at potentials above 30-40 mV. The open state has a weak anion selectivity whereas the closed state is cation-selective (PubMed:11845315, PubMed:18755977, PubMed:20230784, PubMed:8420959). Binds various signaling molecules, including the sphingolipid ceramide, the phospholipid

phosphatidylcholine, and the sterols cholesterol and oxysterol (PubMed:31015432). In depolarized mitochondria, acts downstream of PRKN and PINK1 to promote mitophagy or prevent apoptosis; polyubiquitination by PRKN promotes mitophagy, while monoubiquitination by PRKN decreases mitochondrial calcium influx which ultimately inhibits apoptosis (PubMed:32047033). May participate in the formation of the permeability transition pore complex (PTPC) responsible for the release of mitochondrial products that triggers apoptosis (PubMed:15033708, PubMed:25296756). May mediate ATP export from cells (PubMed:30061676).

Cellular Location

Mitochondrion outer membrane; Multi-pass membrane protein. Cell membrane; Multi-pass membrane protein. Membrane raft; Multi-pass membrane protein

Tissue Location


Expressed in erythrocytes (at protein level) (PubMed:27641616). Expressed in heart, liver and skeletal muscle (PubMed:8420959).

VDAC1 / PORIN Antibody (aa185-197) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

VDAC1 / PORIN Antibody (aa185-197) - Images



Anti-VDAC1 antibody IHC of human liver.

VDAC1 / PORIN Antibody (aa185-197) - Background

Forms a channel through the mitochondrial outer membrane and also the plasma membrane. The

channel at the outer mitochondrial membrane allows diffusion of small hydrophilic molecules; in the plasma membrane it is involved in cell volume regulation and apoptosis. It adopts an open conformation at low or zero membrane potential and a closed conformation at potentials above 30-40 mV. The open state has a weak anion selectivity whereas the closed state is cation-selective. May participate in the formation of the permeability transition pore complex (PTPC) responsible for the release of mitochondrial products that triggers apoptosis.

VDAC1 / PORIN Antibody (aa185-197) - References

- Blachly-Dyson E.,et al.J. Biol. Chem. 268:1835-1841(1993).
Blachly-Dyson E.,et al.Biophys. J. 59:216A-216A(1991).
Decker W.K.,et al.Mamm. Genome 10:1041-1042(1999).
Messina A.,et al.Biochem. Biophys. Res. Commun. 270:787-792(2000).
Ota T.,et al.Nat. Genet. 36:40-45(2004).