

BMF Antibody (C-Terminus)
Rabbit Polyclonal Antibody
Catalog # ALS11390**Specification**

BMF Antibody (C-Terminus) - Product Information

Application	IF, WB, IHC
Primary Accession	O96LC9
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	21kDa KDa

BMF Antibody (C-Terminus) - Additional Information**Gene ID** 90427**Other Names**

Bcl-2-modifying factor, BMF

Target/Specificity

synthetic peptide corresponding to 14 amino acids near the carboxy terminus of human Bmf

Reconstitution & Storage

Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles. Store undiluted.

Precautions

BMF Antibody (C-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

BMF Antibody (C-Terminus) - Protein Information**Name** BMF**Function**

May play a role in apoptosis. Isoform 1 seems to be the main initiator.

Tissue Location

Isoform 1 is mainly expressed in B-lymphoid cells. Isoform 2 and isoform 3 are mainly expressed in B-CLL and normal B- cells.


BMF Antibody (C-Terminus) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


BMF Antibody (C-Terminus) - Images



Immunofluorescence of Bmf in Human Kidney cells with Bmf antibody at 20 ug/ml.



Western blot analysis of Bmf expression in HepG2 cell lysate with Bmf antibody at (A) 2.5 and...



Anti-BMF antibody IHC of human brain, cerebellum.

BMF Antibody (C-Terminus) - Background

May play a role in apoptosis. Isoform 1 seems to be the main initiator.

BMF Antibody (C-Terminus) - References

Puthalakath H., et al. Science 293:1829-1832(2001).
Morales A.A., et al. Leukemia 18:41-47(2004).
Hattori A., et al. DNA Res. 7:357-366(2000).