

Hamartin / TSC1 Antibody
Rabbit Polyclonal Antibody
Catalog # ALS11502**Specification****Hamartin / TSC1 Antibody - Product Information**

Application	ICC, IF, IHC
Primary Accession	Q92574
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	130kDa KDa

Hamartin / TSC1 Antibody - Additional Information**Gene ID** 7248**Other Names**

Hamartin, Tuberous sclerosis 1 protein, TSC1, KIAA0243, TSC

Target/Specificity

15 amino acid peptide from the middle region of human TSC1 TSC1 antibody will recognize both isoforms of TSC1

Reconstitution & Storage

Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles. Store undiluted.

Precautions

Hamartin / TSC1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Hamartin / TSC1 Antibody - Protein Information**Name** TSC1 {ECO:0000303|PubMed:9242607, ECO:0000312|HGNC:HGNC:12362}**Function**

Non-catalytic component of the TSC-TBC complex, a multiprotein complex that acts as a negative regulator of the canonical mTORC1 complex, an evolutionarily conserved central nutrient sensor that stimulates anabolic reactions and macromolecule biosynthesis to promote cellular biomass generation and growth (PubMed: [12172553](http://www.uniprot.org/citations/12172553), PubMed: [12906785](http://www.uniprot.org/citations/12906785), PubMed: [12271141](http://www.uniprot.org/citations/12271141), PubMed: [28215400](http://www.uniprot.org/citations/28215400), PubMed: [15340059](http://www.uniprot.org/citations/15340059), PubMed: [24529379](http://www.uniprot.org/citations/24529379)). The TSC-TBC complex acts as a GTPase-activating protein (GAP) for the small GTPase RHEB, a direct activator of the protein kinase activity of mTORC1 (PubMed: [12906785](http://www.uniprot.org/citations/12906785)).

PubMed:15340059, PubMed:24529379). In absence of nutrients, the TSC-TBC complex inhibits mTORC1, thereby preventing phosphorylation of ribosomal protein S6 kinase (RPS6KB1 and RPS6KB2) and EIF4EBP1 (4E-BP1) by the mTORC1 signaling (PubMed:12271141, PubMed:24529379, PubMed:28215400). The TSC- TBC complex is inactivated in response to nutrients, relieving inhibition of mTORC1 (PubMed:12172553, PubMed:24529379). Within the TSC-TBC complex, TSC1 stabilizes TSC2 and prevents TSC2 self- aggregation (PubMed:10585443, PubMed:28215400). Acts as a tumor suppressor (PubMed:9242607). Involved in microtubule-mediated protein transport via its ability to regulate mTORC1 signaling (By similarity). Also acts as a co-chaperone for HSP90AA1 facilitating HSP90AA1 chaperoning of protein clients such as kinases, TSC2 and glucocorticoid receptor NR3C1 (PubMed:29127155). Increases ATP binding to HSP90AA1 and inhibits HSP90AA1 ATPase activity (PubMed:29127155). Competes with the activating co-chaperone AHSA1 for binding to HSP90AA1, thereby providing a reciprocal regulatory mechanism for chaperoning of client proteins (PubMed:29127155). Recruits TSC2 to HSP90AA1 and stabilizes TSC2 by preventing the interaction between TSC2 and ubiquitin ligase HERC1 (PubMed:16464865, PubMed:29127155).

Cellular Location

Lysosome membrane; Peripheral membrane protein. Cytoplasm, cytosol Note=Recruited to lysosomal membranes in a RHEB-dependent process in absence of nutrients (PubMed:24529379). In response to nutrients, the complex dissociates from lysosomal membranes and relocates to the cytosol (PubMed:24529379).

Tissue Location


Highly expressed in skeletal muscle, followed by heart, brain, placenta, pancreas, lung, liver and kidney (PubMed:9242607). Also expressed in embryonic kidney cells (PubMed:9242607).


Hamartin / TSC1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Hamartin / TSC1 Antibody - Images







Immunocytochemistry of TSC1 in EL4 cells with TSC1 antibody at 2 ug/ml



Immunofluorescence of TSC1 in L1210 cells with TSC1 antibody at 10 ug/ml.



Anti-TSC1 antibody IHC of human skeletal muscle.



Anti-TSC1 antibody IHC of human heart.

Hamartin / TSC1 Antibody - Background

In complex with TSC2, inhibits the nutrient-mediated or growth factor-stimulated phosphorylation of S6K1 and EIF4EBP1 by negatively regulating mTORC1 signaling. Seems not to be required for TSC2 GAP activity towards RHEB. Implicated as a tumor suppressor. Involved in microtubule-mediated protein transport, but this seems to be due to unregulated mTOR signaling.

Hamartin / TSC1 Antibody - References

- van Slegtenhorst M.A.,et al.Science 277:805-808(1997).
- Ota T.,et al.Nat. Genet. 36:40-45(2004).
- Humphray S.J.,et al.Nature 429:369-374(2004).
- Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.
- Nagase T.,et al.DNA Res. 3:321-329(1996).