

HAP1 Antibody (aa639-653)
Rabbit Polyclonal Antibody
Catalog # ALS11579**Specification**

HAP1 Antibody (aa639-653) - Product Information

Application	IHC-P, E
Primary Accession	P54257
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	76kDa KDa
Dilution	IHC-P~~N/A E~~N/A

HAP1 Antibody (aa639-653) - Additional Information**Gene ID** 9001**Other Names**

Huntingtin-associated protein 1, HAP-1, Neuroan 1, HAP1, HAP2, HLP1

Target/Specificity

Amino acids 639 to 653 of human HAP1

Reconstitution & Storage

Long term: -20°C; Short term: +4°C. Avoid repeat freeze-thaw cycles.

Precautions

HAP1 Antibody (aa639-653) is for research use only and not for use in diagnostic or therapeutic procedures.

HAP1 Antibody (aa639-653) - Protein Information**Name** HAP1**Synonyms** HAP2, HLP1**Function**

Originally identified as neuronal protein that specifically associates with HTT/huntingtin and the binding is enhanced by an expanded polyglutamine repeat within HTT possibly affecting HAP1 interaction properties. Both HTT and HAP1 are involved in intracellular trafficking and HAP1 is proposed to link HTT to motor proteins and/or transport cargos. Seems to play a role in vesicular transport within neurons and axons such as from early endosomes to late endocytic compartments and to promote neurite outgrowth. The vesicular transport function via association with microtubule-dependent transporters can be attenuated by association with mutant HTT. Involved in the axonal transport of BDNF and its activity-dependent secretion; the function seems to involve HTT, DCTN1 and a complex with SORT1. Involved in APP trafficking and seems to facilitate APP

anterograde transport and membrane insertion thereby possibly reducing processing into amyloid beta. Involved in delivery of gamma-aminobutyric acid (GABA(A)) receptors to synapses; the function is dependent on kinesin motor protein KIF5 and is disrupted by HTT with expanded polyglutamine repeat. Involved in regulation of autophagosome motility by promoting efficient retrograde axonal transport. Seems to be involved in regulation of membrane receptor recycling and degradation, and respective signal transduction, including GABA(A) receptors, tyrosine kinase receptors, EGFR, IP3 receptor and androgen receptor. Among others suggested to be involved in control of feeding behavior (involving hypothalamic GABA(A) receptors), cerebellar and brainstem development (involving AHI1 and NTRK1/TrkA), postnatal neurogenesis (involving hypothalamic NTRK2/TrkB), and ITPR1/InsP3R1-mediated Ca(2+) release (involving HTT and possibly the effect of mutant HTT). Via association with DCTN1/dynactin p150-glued and HTT/huntingtin involved in cytoplasmic retention of REST in neurons. May be involved in ciliogenesis. Involved in regulation of exocytosis. Seems to be involved in formation of cytoplasmic inclusion bodies (STBs). In case of anomalous expression of TBP, can sequester a subset of TBP into STBs; sequestration is enhanced by an expanded polyglutamine repeat within TBP. HAP1-containing STBs have been proposed to play a protective role against neurodegeneration in Huntington disease (HD) and spinocerebellar ataxia 17 (SCA17).

Cellular Location

Cytoplasm. Cell projection, axon. Presynapse {ECO:0000250|UniProtKB:P54256}. Cytoplasm, cytoskeleton {ECO:0000250|UniProtKB:P54256}. Cell projection, dendritic spine {ECO:0000250|UniProtKB:P54256}. Cell projection, dendrite {ECO:0000250|UniProtKB:P54256}. Lysosome {ECO:0000250|UniProtKB:P54256}. Endoplasmic reticulum {ECO:0000250|UniProtKB:P54256}. Mitochondrion. Nucleus {ECO:0000250|UniProtKB:P54256}. Cytoplasmic vesicle, autophagosome {ECO:0000250|UniProtKB:O35668}. Early endosome {ECO:0000250|UniProtKB:P54256}. Cell projection, growth cone {ECO:0000250|UniProtKB:P54256}. Cell projection, neuron projection {ECO:0000250|UniProtKB:P54256}. Cytoplasmic vesicle, secretory vesicle, synaptic vesicle {ECO:0000250|UniProtKB:P54256}. Note=Localizes to large nonmembrane-bound cytoplasmic bodies found in various types of neurons, called stigmoid bodies (STBs). Localization to neuronal processes and neurite tips is decreased by YWHAZ. In the nucleus localizes to nuclear rods. {ECO:0000250|UniProtKB:P54256}

Tissue Location

Predominantly expressed in brain. Selectively expressed in neurons


HAP1 Antibody (aa639-653) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

HAP1 Antibody (aa639-653) - Images





Anti-HAP1 antibody IHC of human adrenal.

HAP1 Antibody (aa639-653) - Background

Originally identified as neuronal protein that specifically associates with HTT/huntingtin and the binding is enhanced by an expanded polyglutamine repeat within HTT possibly affecting HAP1 interaction properties. Both HTT and HAP1 are involved in intracellular trafficking and HAP1 is proposed to link HTT to motor proteins and/or transport cargos. Seems to play a role in vesicular transport within neurons and axons such as from early endosomes to late endocytic compartments and to promote neurite outgrowth. The vesicular transport function via association with microtubule-dependent transporters can be attenuated by association with mutant HTT. Involved in the axonal transport of BDNF and its activity-dependent secretion; the function seems to involve HTT, DCTN1 and a complex with SORT1. Involved in APP trafficking and seems to facilitate APP anterograde transport and membrane insertion thereby possibly reducing processing into amyloid beta. Involved in delivery of gamma-aminobutyric acid (GABA(A)) receptors to synapses; the function is dependent on kinesin motor protein KIF5 and is disrupted by HTT with expanded polyglutamine repeat. Involved in regulation of autophagosome motility by promoting efficient retrograde axonal transport. Seems to be involved in regulation of membrane receptor recycling and degradation, and respective signal transduction, including GABA(A) receptors, tyrosine kinase receptors, EGFR, IP3 receptor and androgen receptor. Among others suggested to be involved in control of feeding behavior (involving hypothalamic GABA(A) receptors), cerebellar and brainstem development (involving AHI1 and NTRK1/TrkA), postnatal neurogenesis (involving hypothalamic NTRK2/TrkB), and ITPR1/InsP3R1-mediated Ca(2+) release (involving HTT and possibly the effect of mutant HTT). Via association with DCTN1/dynactin p150-glued and HTT/huntingtin involved in cytoplasmic retention of REST in neurons. May be involved in ciliogenesis. Involved in regulation of exocytosis. Seems to be involved in formation of cytoplasmic inclusion bodies (STBs). In case of anomalous expression of TBP, can sequester a subset of TBP into STBs; sequestration is enhanced by an expanded polyglutamine repeat within TBP. HAP1-containing STBs have been proposed to play a protective role against neurodegeneration in Huntington disease (HD) and spinocerebellar ataxia 17 (SCA17).

HAP1 Antibody (aa639-653) - References

- Li S.-H., et al. J. Biol. Chem. 273:19220-19227(1998).
- Nasir J., et al. Gene 254:181-187(2000).
- Ota T., et al. Nat. Genet. 36:40-45(2004).
- Zody M.C., et al. Nature 440:1045-1049(2006).
- Totoki Y., et al. Submitted (MAR-2005) to the EMBL/GenBank/DBJ databases.