

# **SPHK2 Antibody (N-Terminus)**

Rabbit Polyclonal Antibody Catalog # ALS11619

# **Specification**

# SPHK2 Antibody (N-Terminus) - Product Information

Application
Primary Accession
Reactivity
Host
Clonality
Calculated MW
Dilution

WB, IHC-P, E
O9NRA0
Human
Rabbit
Polyclonal
69kDa KDa
WB~~1:1000
IHC-P~~N/A
E~~N/A

# SPHK2 Antibody (N-Terminus) - Additional Information

**Gene ID** 56848

### **Other Names**

Sphingosine kinase 2, SK 2, SPK 2, 2.7.1.91, SPHK2

### Target/Specificity

Peptide from the N-terminus of human SPHK2 (long form)

### **Reconstitution & Storage**

+4°C or -20°C, Avoid repeated freezing and thawing.

#### **Precautions**

SPHK2 Antibody (N-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

# SPHK2 Antibody (N-Terminus) - Protein Information

Name SPHK2 (<u>HGNC:18859</u>)

Synonyms SK2

## **Function**

Catalyzes the phosphorylation of sphingosine to form sphingosine-1-phosphate (SPP), a lipid mediator with both intra- and extracellular functions. Also acts on D-erythro-dihydrosphingosine, D- erythro-sphingosine and L-threo-dihydrosphingosine. Binds phosphoinositides (PubMed:<a href="http://www.uniprot.org/citations/12954646" target="\_blank">12954646</a>, PubMed:<a href="http://www.uniprot.org/citations/19168031" target="\_blank">19168031</a>). In contrast to prosurvival SPHK1, has a positive effect on intracellular ceramide levels, inhibits cells growth and enhances apoptosis (PubMed:<a href="http://www.uniprot.org/citations/16118219" target="\_blank">16118219</a></a>). In mitochondria, is important for cytochrome-c oxidase



assembly and mitochondrial respiration. The SPP produced in mitochondria binds PHB2 and modulates the regulation via PHB2 of complex IV assembly and respiration (PubMed:<a href="http://www.uniprot.org/citations/20959514" target="\_blank">20959514</a>). In nucleus, plays a role in epigenetic regulation of gene expression. Interacts with HDAC1 and HDAC2 and, through SPP production, inhibits their enzymatic activity, preventing the removal of acetyl groups from lysine residues with histones. Up- regulates acetylation of histone H3-K9, histone H4-K5 and histone H2B- K12 (PubMed:<a href="http://www.uniprot.org/citations/19729656" target="\_blank">19729656</a>). In nucleus, may have an inhibitory effect on DNA synthesis and cell cycle (PubMed:<a href="http://www.uniprot.org/citations/12954646" target="\_blank">12954646</a>, PubMed:<a href="http://www.uniprot.org/citations/12954646" target="\_blank">16103110</a> (a>). In mast cells, is the main regulator of SPP production which mediates calcium influx, NF-kappa-B activation, cytokine production, such as TNF and IL6, and degranulation of mast cells (By similarity). In dopaminergic neurons, is involved in promoting mitochondrial functions regulating ATP and ROS levels (By similarity). Also involved in the regulation of glucose and lipid metabolism (By similarity).

### **Cellular Location**

Cytoplasm. Nucleus. Endoplasmic reticulum {ECO:0000250|UniProtKB:Q9JIA7}. Mitochondrion inner membrane {ECO:0000250|UniProtKB:Q9JIA7}. Note=In nucleus, located in nucleosomes where it associates with core histone proteins such as histone 3 (PubMed:19729656). In brains of patients with Alzheimer's disease, may be preferentially localized in the nucleus. Cytosolic expression decrease correlates with the density of amyloid deposits (PubMed:29615132). In apoptotic cells, colocalizes with CASP1 in cell membrane where is cleaved and released from cells in an active form (PubMed:20197547).

#### **Tissue Location**

Mainly expressed in adult kidney, liver, and brain (PubMed:10751414). Expressed in cerebral cortex and hippocampus (at protein level) (PubMed:29615132). Isoform 1 is the predominant form expressed in most tissues (PubMed:16103110)


# SPHK2 Antibody (N-Terminus) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

# SPHK2 Antibody (N-Terminus) - Images





Anti-SPHK2 antibody IHC of human brain, cerebellum.

# SPHK2 Antibody (N-Terminus) - Background

Catalyzes the phosphorylation of sphingosine to form sphingosine 1-phosphate (SPP), a lipid mediator with both intra- and extracellular functions. Also acts on D-erythro- dihydrosphingosine, D-erythro-sphingosine and L-threo- dihydrosphingosine. Binds phosphoinositides.

# **SPHK2 Antibody (N-Terminus) - References**

Liu H.,et al.J. Biol. Chem. 275:19513-19520(2000). Alemany R.,et al.Submitted (NOV-2006) to the EMBL/GenBank/DDBJ databases. Wiemann S.,et al.Genome Res. 11:422-435(2001). Ota T.,et al.Nat. Genet. 36:40-45(2004). Grimwood J.,et al.Nature 428:529-535(2004).