

MICA Antibody (C-Terminus)
Rabbit Polyclonal Antibody
Catalog # ALS11697**Specification**

MICA Antibody (C-Terminus) - Product Information

Application	IHC
Primary Accession	Q29983
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	43kDa KDa

MICA Antibody (C-Terminus) - Additional Information**Gene ID** 100507436**Other Names**

MHC class I polypeptide-related sequence A, MIC-A, MICA {ECO:0000312|EMBL:CAI41907.1}

Target/Specificity

a 16 amino acid peptide from near the carboxy terminus of human MICA

Reconstitution & Storage

Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles. Store undiluted.

Precautions

MICA Antibody (C-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

MICA Antibody (C-Terminus) - Protein Information**Name** MICA {ECO:0000312|EMBL:CAI41907.1}**Function**

Seems to have no role in antigen presentation. Acts as a stress-induced self-antigen that is recognized by gamma delta T-cells. Ligand for the KLRK1/NKG2D receptor. Binding to KLRK1 leads to cell lysis.

Cellular Location

Cell membrane; Single-pass type I membrane protein. Cytoplasm. Note=Expressed on the cell surface in gastric epithelium, endothelial cells and fibroblasts and in the cytoplasm in keratinocytes and monocytes. Infection with human adenovirus 5 suppresses cell surface expression due to the adenoviral E3-19K protein which causes retention in the endoplasmic reticulum

Tissue Location

Widely expressed with the exception of the central nervous system where it is absent. Expressed


predominantly in gastric epithelium and also in monocytes, keratinocytes, endothelial cells, fibroblasts and in the outer layer of Hassal's corpuscles within the medulla of normal thymus. In skin, expressed mainly in the keratin layers, basal cells, ducts and follicles. Also expressed in many, but not all, epithelial tumors of lung, breast, kidney, ovary, prostate and colon. In thymomas, overexpressed in cortical and medullar epithelial cells. Tumors expressing MICA display increased levels of gamma delta T-cells.

MICA Antibody (C-Terminus) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MICA Antibody (C-Terminus) - Images



Anti-MICA antibody IHC of human prostate.

MICA Antibody (C-Terminus) - Background

Seems to have no role in antigen presentation. Acts as a stress-induced self-antigen that is recognized by gamma delta T- cells. Ligand for the KLRK1/NKG2D receptor. Binding to KLRK1 leads to cell lysis.

MICA Antibody (C-Terminus) - References

Bahram S.,et al.Proc. Natl. Acad. Sci. U.S.A. 91:6259-6263(1994).
Bahram S.,et al.Immunogenetics 44:80-81(1996).
Mungall A.J.,et al.Nature 425:805-811(2003).
Leelayuwat C.,et al.Immunogenetics 40:339-351(1994).
Fodil N.,et al.Immunogenetics 44:351-357(1996).