

VISA / MAVS Antibody (Internal)
Rabbit Polyclonal Antibody
Catalog # ALS11742**Specification****VISA / MAVS Antibody (Internal) - Product Information**

Application	WB, IHC-P
Primary Accession	Q7Z434
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	57kDa KDa
Dilution	WB~~1:1000 IHC-P~~N/A

VISA / MAVS Antibody (Internal) - Additional Information**Gene ID** 57506**Other Names**

Mitochondrial antiviral-signaling protein, MAVS, CARD adapter inducing interferon beta, Cardif, Interferon beta promoter stimulator protein 1, IPS-1, Putative NF-kappa-B-activating protein 031N, Virus-induced-signaling adapter, VISA, MAVS, IPS1, KIAA1271, VISA

Target/Specificity

17 amino acid peptide from near the center of human VISA.

Reconstitution & Storage

Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles. Store undiluted.

Precautions

VISA / MAVS Antibody (Internal) is for research use only and not for use in diagnostic or therapeutic procedures.

VISA / MAVS Antibody (Internal) - Protein Information**Name** MAVS {ECO:0000303|PubMed:16125763, ECO:0000312|HGNC:HGNC:29233}**Function**

Adapter required for innate immune defense against viruses (PubMed:16125763, PubMed:16127453, PubMed:16153868, PubMed:16177806, PubMed:19631370, PubMed:20127681, PubMed:20451243, PubMed:21170385, PubMed:<a

[23087404](http://www.uniprot.org/citations/23087404), PubMed: [27992402](http://www.uniprot.org/citations/27992402), PubMed: [33139700](http://www.uniprot.org/citations/33139700), PubMed: [37582970](http://www.uniprot.org/citations/37582970)). Acts downstream of DHX33, RIGI and IFIH1/MDA5, which detect intracellular dsRNA produced during viral replication, to coordinate pathways leading to the activation of NF-kappa-B, IRF3 and IRF7, and to the subsequent induction of antiviral cytokines such as IFNB and RANTES (CCL5) (PubMed: [16125763](http://www.uniprot.org/citations/16125763), PubMed: [16127453](http://www.uniprot.org/citations/16127453), PubMed: [16153868](http://www.uniprot.org/citations/16153868), PubMed: [16177806](http://www.uniprot.org/citations/16177806), PubMed: [19631370](http://www.uniprot.org/citations/19631370), PubMed: [20127681](http://www.uniprot.org/citations/20127681), PubMed: [20451243](http://www.uniprot.org/citations/20451243), PubMed: [20628368](http://www.uniprot.org/citations/20628368), PubMed: [21170385](http://www.uniprot.org/citations/21170385), PubMed: [23087404](http://www.uniprot.org/citations/23087404), PubMed: [25636800](http://www.uniprot.org/citations/25636800), PubMed: [27736772](http://www.uniprot.org/citations/27736772), PubMed: [33110251](http://www.uniprot.org/citations/33110251)). Peroxisomal and mitochondrial MAVS act sequentially to create an antiviral cellular state (PubMed: [20451243](http://www.uniprot.org/citations/20451243)). Upon viral infection, peroxisomal MAVS induces the rapid interferon-independent expression of defense factors that provide short-term protection, whereas mitochondrial MAVS activates an interferon-dependent signaling pathway with delayed kinetics, which amplifies and stabilizes the antiviral response (PubMed: [20451243](http://www.uniprot.org/citations/20451243)). May activate the same pathways following detection of extracellular dsRNA by TLR3 (PubMed: [16153868](http://www.uniprot.org/citations/16153868)). May protect cells from apoptosis (PubMed: [16125763](http://www.uniprot.org/citations/16125763)). Involved in NLRP3 inflammasome activation by mediating NLRP3 recruitment to mitochondria (PubMed: [23582325](http://www.uniprot.org/citations/23582325)).

Cellular Location

Mitochondrion outer membrane; Single-pass membrane protein. Mitochondrion. Peroxisome

Tissue Location


Present in T-cells, monocytes, epithelial cells and hepatocytes (at protein level). Ubiquitously expressed, with highest levels in heart, skeletal muscle, liver, placenta and peripheral blood leukocytes.

VISA / MAVS Antibody (Internal) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

VISA / MAVS Antibody (Internal) - Images



Anti-MAVS antibody IHC of human brain, cortex.

VISA / MAVS Antibody (Internal) - Background

Required for innate immune defense against viruses. Acts downstream of DDX58/RIG-I and IFIH1/MDA5, which detect intracellular dsRNA produced during viral replication, to coordinate pathways leading to the activation of NF-kappa-B, IRF3 and IRF7, and to the subsequent induction of antiviral cytokines such as IFN-beta and RANTES (CCL5). Peroxisomal and mitochondrial MAVS act sequentially to create an antiviral cellular state. Upon viral infection, peroxisomal MAVS induces the rapid interferon- independent expression of defense factors that provide short-term protection, whereas mitochondrial MAVS activates an interferon- dependent signaling pathway with delayed kinetics, which amplifies and stabilizes the antiviral response. May activate the same pathways following detection of extracellular dsRNA by TLR3. May protect cells from apoptosis.

VISA / MAVS Antibody (Internal) - References

- Seth R.B.,et al.Cell 122:669-682(2005).
- Xu L.-G.,et al.Mol. Cell 19:727-740(2005).
- Meylan E.,et al.Nature 437:1167-1172(2005).
- Kawai T.,et al.Nat. Immunol. 6:981-988(2005).
- Lad S.P.,et al.Mol. Immunol. 45:2277-2287(2008).