

STMN1 / Stathmin / LAG Antibody (Ser37)
Rabbit Polyclonal Antibody
Catalog # ALS11780**Specification**

STMN1 / Stathmin / LAG Antibody (Ser37) - Product Information

Application	WB, IF, IHC
Primary Accession	P16949
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	17kDa KDa

STMN1 / Stathmin / LAG Antibody (Ser37) - Additional Information**Gene ID** 3925**Other Names**

Stathmin, Leukemia-associated phosphoprotein p18, Metablastin, Oncoprotein 18, Op18, Phosphoprotein p19, pp19, Prosolin, Protein Pr22, pp17, STMN1, C1orf215, LAP18, OP18

Target/Specificity

Amino acids surrounding Ser 37 of human STMN1

Reconstitution & Storage

Long term: -70°C; Short term: -20°C

Precautions

STMN1 / Stathmin / LAG Antibody (Ser37) is for research use only and not for use in diagnostic or therapeutic procedures.

STMN1 / Stathmin / LAG Antibody (Ser37) - Protein Information**Name** STMN1**Synonyms** C1orf215, LAP18, OP18**Function**

Involved in the regulation of the microtubule (MT) filament system by destabilizing microtubules. Prevents assembly and promotes disassembly of microtubules. Phosphorylation at Ser-16 may be required for axon formation during neurogenesis. Involved in the control of the learned and innate fear (By similarity).

Cellular Location

Cytoplasm, cytoskeleton.

Tissue Location

Ubiquitous. Expression is strongest in fetal and adult brain, spinal cord, and cerebellum, followed


by thymus, bone marrow, testis, and fetal liver. Expression is intermediate in colon, ovary, placenta, uterus, and trachea, and is readily detected at substantially lower levels in all other tissues examined. Lowest expression is found in adult liver. Present in much greater abundance in cells from patients with acute leukemia of different subtypes than in normal peripheral blood lymphocytes, non-leukemic proliferating lymphoid cells, bone marrow cells, or cells from patients with chronic lymphoid or myeloid leukemia.

STMN1 / Stathmin / LAG Antibody (Ser37) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


STMN1 / Stathmin / LAG Antibody (Ser37) - Images



Western blot of extracts from 293 and HeLa cells using Stathmin 1(Ab-38) antibody.



Immunofluorescence staining of methanol-fixed HeLa cells using Stathmin 1(Ab-38) antibody.



Anti-STMN1 antibody IHC of human brain, cerebellum.

STMN1 / Stathmin / LAG Antibody (Ser37) - Background

Involved in the regulation of the microtubule (MT) filament system by destabilizing microtubules. Prevents assembly and promotes disassembly of microtubules. Phosphorylation at Ser- 16 may be required for axon formation during neurogenesis. Involved in the control of the learned and innate fear (By similarity).

STMN1 / Stathmin / LAG Antibody (Ser37) - References

Zhu X.-X.,et al.J. Biol. Chem. 264:14556-14560(1989).
Maucuer A.,et al.FEBS Lett. 264:275-278(1990).
Melhem R.F.,et al.J. Biol. Chem. 266:17747-17753(1991).
Hosoya H.,et al.Cell Struct. Funct. 21:237-243(1996).
Ota T.,et al.Nat. Genet. 36:40-45(2004).