10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

IKBKG / NEMO / IKK Gamma Antibody (aa2-13)

Rabbit Polyclonal Antibody
Catalog # ALS11881

Specification

IKBKG / NEMO / IKK Gamma Antibody (aa2-13) - Product Information

Application	**WB, IHC-P, E**
Primary Accession	Q9Y6K9
Reactivity	**Human**
Host	**Rabbit**
Clonality	**Polyclonal**
Calculated MW	**48kDa KDa**
Dilution	**WB~~1:1000** **IHC-P~~N/A** **E~~N/A**

IKBKG / NEMO / IKK Gamma Antibody (aa2-13) - Additional Information

Gene ID 8517

Other Names
NF-kappa-B essential modulator, NEMO, FIP-3, IkB kinase-associated protein 1, IKKAP1, Inhibitor of nuclear factor kappa-B kinase subunit gamma, I-kappa-B kinase subunit gamma, IKK-gamma, IKKG, IkB kinase subunit gamma, NF-kappa-B essential modifier, IKBKG, FIP3, NEMO

Target/Specificity
Residues 2-13 [NRHLWKSQLCEM] of the IKBKG protein

Reconstitution & Storage
+4°C or -20°C, Avoid repeated freezing and thawing.

Precautions
IKBKG / NEMO / IKK Gamma Antibody (aa2-13) is for research use only and not for use in diagnostic or therapeutic procedures.

IKBKG / NEMO / IKK Gamma Antibody (aa2-13) - Protein Information

Name IKBKG (HGNC:5961)

Synonyms FIP3, NEMO

Function
Regulatory subunit of the IKK core complex which phosphorylates inhibitors of NF-kappa-B thus leading to the dissociation of the inhibitor/NF-kappa-B complex and ultimately the degradation of the inhibitor (PubMed:14695475, PubMed:20724660, PubMed:21518757, PubMed:<a href="http://www.uniprot.org/citations/9751060"

target="_blank">9751060). Its binding to scaffolding polyubiquitin plays a key role in IKK activation by multiple signaling receptor pathways (PubMed:16547522, PubMed:18287044, PubMed:19033441, PubMed:19185524, PubMed:21606507, PubMed:27777308, PubMed:33567255). Can recognize and bind both 'Lys-63'-linked and linear polyubiquitin upon cell stimulation, with a much higher affinity for linear polyubiquitin (PubMed:16547522, PubMed:18287044, PubMed:19033441, PubMed:19185524, PubMed:21606507, PubMed:27777308). Could be implicated in NF-kappa-B-mediated protection from cytokine toxicity. Essential for viral activation of IRF3 (PubMed:19854139). Involved in TLR3- and IFIH1-mediated antiviral innate response; this function requires 'Lys- 27'-linked polyubiquitination (PubMed:20724660).

Cellular Location
Cytoplasm. Nucleus Note=Sumoylated NEMO accumulates in the nucleus in response to genotoxic stress.

Tissue Location
Heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas

IKBKG / NEMO / IKK Gamma Antibody (aa2-13) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

IKBKG / NEMO / IKK Gamma Antibody (aa2-13) - Images

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999



Anti-IKBKG / IKK Gamma antibody IHC of human kidney.

IKBKG / NEMO / IKK Gamma Antibody (aa2-13) - Background

Regulatory subunit of the IKK core complex which phosphorylates inhibitors of NF-kappa-B thus leading to the dissociation of the inhibitor/NF-kappa-B complex and ultimately the degradation of the inhibitor. Its binding to scaffolding polyubiquitin seems to play a role in IKK activation by multiple signaling receptor pathways. However, the specific type of polyubiquitin recognized upon cell stimulation (either 'Lys-63'- linked or linear polyubiquitin) and its functional importance is reported conflictingly. Also considered to be a mediator for TAX activation of NF-kappa-B. Could be implicated in NF-kappa-B- mediated protection from cytokine toxicity. Essential for viral activation of IRF3. Involved in TLR3- and IFIH1-mediated antiviral innate response; this function requires 'Lys-27'-linked polyubiquitination.

IKBKG / NEMO / IKK Gamma Antibody (aa2-13) - References

Li Y.,et al.Proc. Natl. Acad. Sci. U.S.A. 96:1042-1047(1999).
Jin D.-Y.,et al.J. Biomed. Sci. 6:115-120(1999).
Rothwarf D.M.,et al.Nature 395:297-300(1998).
Smahi A.,et al.Nature 405:466-472(2000).
Ye Z.,et al.Biochem. Biophys. Res. Commun. 275:223-227(2000).