

HCAR3 / GPR109B / HM74 Antibody (aa1-50)
Rabbit Polyclonal Antibody
Catalog # ALS11914**Specification**

HCAR3 / GPR109B / HM74 Antibody (aa1-50) - Product Information

Application	WB
Primary Accession	P49019
Reactivity	Human, Hamster, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	44kDa KDa

HCAR3 / GPR109B / HM74 Antibody (aa1-50) - Additional Information**Gene ID** 8843**Other Names**

Hydroxycarboxylic acid receptor 3, G-protein coupled receptor 109B, G-protein coupled receptor HM74, G-protein coupled receptor HM74B, Niacin receptor 2, Nicotinic acid receptor 2, HCAR3, GPR109B, HCA3, HM74B, NIACR2

Target/Specificity

A portion of amino acids 1-50 of human GPR109B

Reconstitution & Storage

Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles.

Precautions

HCAR3 / GPR109B / HM74 Antibody (aa1-50) is for research use only and not for use in diagnostic or therapeutic procedures.

HCAR3 / GPR109B / HM74 Antibody (aa1-50) - Protein Information**Name** HCAR3**Synonyms** GPR109B, HCA3, HM74B, NIACR2**Function**

Receptor for 3-OH-octanoid acid mediates a negative feedback regulation of adipocyte lipolysis to counteract prolipolytic influences under conditions of physiological or pathological increases in beta- oxidation rates. Acts as a low affinity receptor for nicotinic acid. This pharmacological effect requires nicotinic acid doses that are much higher than those provided by a normal diet.

Cellular Location

Cell membrane; Multi-pass membrane protein.

Tissue Location


Expression largely restricted to adipose tissue and spleen.

HCAR3 / GPR109B / HM74 Antibody (aa1-50) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

HCAR3 / GPR109B / HM74 Antibody (aa1-50) - Images



Western blot of human GPR109B in human liver lysate using antibody at 1 ug/ml.

HCAR3 / GPR109B / HM74 Antibody (aa1-50) - Background

Receptor for 3-OH-octanoid acid mediates a negative feedback regulation of adipocyte lipolysis to counteract prolipolytic influences under conditions of physiological or pathological increases in beta-oxidation rates. Acts as a low affinity receptor for nicotinic acid. This pharmacological effect requires nicotinic acid doses that are much higher than those provided by a normal diet.

HCAR3 / GPR109B / HM74 Antibody (aa1-50) - References

Nomura H.,et al.Int. Immunol. 5:1239-1249(1993).
Suwa M.,et al.Submitted (JUL-2001) to the EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Scherer S.E.,et al.Nature 440:346-351(2006).
Wise A.,et al.J. Biol. Chem. 278:9869-9874(2003).