

PCNA / Cyclin Antibody (clone PC10)

Mouse Monoclonal Antibody Catalog # ALS11995

## Specification

## PCNA / Cyclin Antibody (clone PC10) - Product Information

Application Primary Accession Reactivity Host Clonality Calculated MW Dilution WB, IHC-P, IP, FC <u>P12004</u> Human, Mouse Monoclonal 29kDa KDa WB~~1:1000 IHC-P~~N/A IP~~N/A FC~~1:10~50

### PCNA / Cyclin Antibody (clone PC10) - Additional Information

Gene ID 5111

Other Names Proliferating cell nuclear antigen, PCNA, Cyclin, PCNA

Target/Specificity Recombinant rat PCNA

**Reconstitution & Storage** Store at 4°C. Do not freeze.

**Precautions** PCNA / Cyclin Antibody (clone PC10) is for research use only and not for use in diagnostic or therapeutic procedures.

### PCNA / Cyclin Antibody (clone PC10) - Protein Information

#### Name PCNA

#### Function

Auxiliary protein of DNA polymerase delta and epsilon, is involved in the control of eukaryotic DNA replication by increasing the polymerase's processibility during elongation of the leading strand (PubMed:<a href="http://www.uniprot.org/citations/35585232" target="\_blank">35585232</a>). Induces a robust stimulatory effect on the 3'-5' exonuclease and 3'-phosphodiesterase, but not apurinic-apyrimidinic (AP) endonuclease, APEX2 activities. Has to be loaded onto DNA in order to be able to stimulate APEX2. Plays a key role in DNA damage response (DDR) by being conveniently positioned at the replication fork to coordinate DNA replication with DNA repair and DNA damage tolerance pathways (PubMed:<a href="http://www.uniprot.org/citations/24939902" target="\_blank">>24939902</a>). Acts as a loading platform to recruit DDR proteins that allow



completion of DNA replication after DNA damage and promote postreplication repair: Monoubiquitinated PCNA leads to recruitment of translesion (TLS) polymerases, while 'Lys-63'-linked polyubiquitination of PCNA is involved in error-free pathway and employs recombination mechanisms to synthesize across the lesion (PubMed:<a href="http://www.uniprot.org/citations/24695737" target=" blank">>24695737</a>).

#### **Cellular Location**

Nucleus. Note=Colocalizes with CREBBP, EP300 and POLD1 to sites of DNA damage (PubMed:24939902). Forms nuclear foci representing sites of ongoing DNA replication and vary in morphology and number during S phase (PubMed:15543136). Co-localizes with SMARCA5/SNF2H and BAZ1B/WSTF at replication foci during S phase (PubMed:15543136). Together with APEX2, is redistributed in discrete nuclear foci in presence of oxidative DNA damaging agents

Volume 50 μl

### PCNA / Cyclin Antibody (clone PC10) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- <u>Dot Blot</u>
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

PCNA / Cyclin Antibody (clone PC10) - Images



Anti-PCNA antibody IHC of human placenta.





Hela cell nuclear extract was resolved by electrophoresis, transferred to nitrocellulose and...

# PCNA / Cyclin Antibody (clone PC10) - Background

Auxiliary protein of DNA polymerase delta and is involved in the control of eukaryotic DNA replication by increasing the polymerase's processibility during elongation of the leading strand. Induces a robust stimulatory effect on the 3'- 5' exonuclease and 3'-phosphodiesterase, but not apurinic- apyrimidinic (AP) endonuclease, APEX2 activities. Has to be loaded onto DNA in order to be able to stimulate APEX2. Plays a key role in DNA damage response (DDR) by being conveniently positioned at the replication fork to coordinate DNA replication with DNA repair and DNA damage tolerance pathways. Acts as a loading platform to recruit DDR proteins that allow completion of DNA replication after DNA damage and promote postreplication repair: Monoubiquitinated PCNA leads to recruitment of translesion (TLS) polymerases, while 'Lys-63'-linked polyubiquitination of PCNA is involved in error-free pathway and employs recombination mechanisms to synthesize across the lesion.

# PCNA / Cyclin Antibody (clone PC10) - References

Almendral J.M., et al. Proc. Natl. Acad. Sci. U.S.A. 84:1575-1579(1987). Travali S., et al.J. Biol. Chem. 264:7466-7472(1989). Ota T., et al.Nat. Genet. 36:40-45(2004). Deloukas P., et al.Nature 414:865-871(2001). Mural R.J., et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.