

CD38 Antibody (clone HIT2)
Mouse Monoclonal Antibody
Catalog # ALS12024**Specification**

CD38 Antibody (clone HIT2) - Product Information

Application	IHC
Primary Accession	P28907
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Calculated MW	34kDa KDa

CD38 Antibody (clone HIT2) - Additional Information**Gene ID** 952**Other Names**

ADP-ribosyl cyclase/cyclic ADP-ribose hydrolase 1, 3.2.2.6, 2'-phospho-ADP-ribosyl cyclase, 2'-phospho-ADP-ribosyl cyclase/2'-phospho-cyclic-ADP-ribose transferase, 2.4.99.20, 2'-phospho-cyclic-ADP-ribose transferase, ADP-ribosyl cyclase 1, ADPRC 1, Cyclic ADP-ribose hydrolase 1, cADPr hydrolase 1, T10, CD38, CD38

Reconstitution & Storage

Store undiluted at 4 degrees C

Precautions

CD38 Antibody (clone HIT2) is for research use only and not for use in diagnostic or therapeutic procedures.

CD38 Antibody (clone HIT2) - Protein Information**Name** CD38**Function**

Synthesizes cyclic ADP-ribose (cADPR), a second messenger for glucose-induced insulin secretion (PubMed:8253715, PubMed:7961800). Synthesizes the Ca(2+) mobilizer nicotinate-adenine dinucleotide phosphate, NAADP(+), from 2'-phospho-cADPR and nicotinic acid, as well as from NADP(+) and nicotinic acid. At both pH 5.0 and pH 7.4 preferentially transforms 2'-phospho-cADPR into NAADP(+), while preferentially cleaving NADP(+) to cADPR and ADPRP rather than into NADDP(+) (PubMed:16690024). Has cADPR hydrolase activity (PubMed:8253715, PubMed:7961800).

Cellular Location

Cell surface. Membrane; Single-pass type II membrane protein


Tissue Location

Expressed at high levels in pancreas, liver, kidney, brain, testis, ovary, placenta, malignant lymphoma and neuroblastoma.

CD38 Antibody (clone HIT2) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CD38 Antibody (clone HIT2) - Images

Anti-CD38 antibody IHC of human prostate.

CD38 Antibody (clone HIT2) - Background

Synthesizes the second messengers cyclic ADP-ribose and nicotinate-adenine dinucleotide phosphate, the former a second messenger for glucose-induced insulin secretion. Also has cADPr hydrolase activity. Also moonlights as a receptor in cells of the immune system.

CD38 Antibody (clone HIT2) - References

Jackson D.G.,et al.J. Immunol. 144:2811-2815(1990).
Nata K.,et al.Gene 186:285-292(1997).
States D.J.,et al.Trends Biochem. Sci. 17:495-495(1992).
Takasawa S.,et al.J. Biol. Chem. 268:26052-26054(1993).
Tohgo A.,et al.J. Biol. Chem. 269:28555-28557(1994).