

**PRMT7 Antibody (aa346-360)**  
**Rabbit Polyclonal Antibody**  
**Catalog # ALS12042****Specification**

---

**PRMT7 Antibody (aa346-360) - Product Information**

Application	WB
Primary Accession	<a href="#">O9NVM4</a>
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	78kDa KDa

**PRMT7 Antibody (aa346-360) - Additional Information****Gene ID** 54496**Other Names**

Protein arginine N-methyltransferase 7, 2.1.1.-, Histone-arginine N-methyltransferase PRMT7, 2.1.1.125, [Myelin basic protein]-arginine N-methyltransferase PRMT7, 2.1.1.126, PRMT7, KIAA1933

**Target/Specificity**

Amino acids 346-360 of human PRMT7 were used as the immunogen.

**Reconstitution & Storage**

Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles.

**Precautions**

PRMT7 Antibody (aa346-360) is for research use only and not for use in diagnostic or therapeutic procedures.

**PRMT7 Antibody (aa346-360) - Protein Information****Name** PRMT7**Synonyms** KIAA1933**Function**

Arginine methyltransferase that can both catalyze the formation of omega-N monomethylarginine (MMA) and symmetrical dimethylarginine (sDMA), with a preference for the formation of MMA. Specifically mediates the symmetrical dimethylation of arginine residues in the small nuclear ribonucleoproteins Sm D1 (SNRPD1) and Sm D3 (SNRPD3); such methylation being required for the assembly and biogenesis of snRNP core particles. Specifically mediates the symmetric dimethylation of histone H4 'Arg-3' to form H4R3me2s. Plays a role in gene imprinting by being recruited by CTCFL at the H19 imprinted control region (ICR) and methylating histone H4 to form H4R3me2s, possibly leading to recruit DNA methyltransferases at these sites. May also play a role in embryonic stem cell (ESC) pluripotency. Also able to mediate the arginine methylation of

histone H2A and myelin basic protein (MBP) in vitro; the relevance of such results is however unclear in vivo.

**Cellular Location**

Cytoplasm, cytosol. Nucleus


**Volume**

50 µl

**PRMT7 Antibody (aa346-360) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**PRMT7 Antibody (aa346-360) - Images**

Western blot of PRMT7 in (A) HeLa and (B) NIH 3T3 cell lysate using antibody at 0.5 µg/ml.

**PRMT7 Antibody (aa346-360) - Background**

Arginine methyltransferase that can both catalyze the formation of omega-N monomethylarginine (MMA) and symmetrical dimethylarginine (sDMA), with a preference for the formation of MMA. Specifically mediates the symmetrical dimethylation of arginine residues in the small nuclear ribonucleoproteins Sm D1 (SNRPD1) and Sm D3 (SNRPD3); such methylation being required for the assembly and biogenesis of snRNP core particles. Specifically mediates the symmetric dimethylation of histone H4 'Arg-3' to form H4R3me2s. Plays a role in gene imprinting by being recruited by CTCFL at the H19 imprinted control region (ICR) and methylating histone H4 to form H4R3me2s, possibly leading to recruit DNA methyltransferases at these sites. May also play a role in embryonic stem cell (ESC) pluripotency. Also able to mediate the arginine methylation of histone H2A and myelin basic protein (MBP) in vitro; the relevance of such results is however unclear in vivo.

**PRMT7 Antibody (aa346-360) - References**

Ota T.,et al.Nat. Genet. 36:40-45(2004).  
Nagase T.,et al.DNA Res. 8:179-187(2001).  
Miranda T.B.,et al.J. Biol. Chem. 279:22902-22907(2004).  
Lee J.-H.,et al.J. Biol. Chem. 280:3656-3664(2005).  
Gonsalvez G.B.,et al.J. Cell Biol. 178:733-740(2007).