

LF / LTF / Lactoferrin Antibody (Internal)

Goat Polyclonal Antibody Catalog # ALS12620

Specification

LF / LTF / Lactoferrin Antibody (Internal) - Product Information

Application WB, IHC
Primary Accession P02788
Reactivity Human
Host Goat
Clonality Polyclonal
Calculated MW 78kDa KDa

LF / LTF / Lactoferrin Antibody (Internal) - Additional Information

Gene ID 4057

Other Names

Lactotransferrin, Lactoferrin, 3.4.21.-, Growth-inhibiting protein 12, Talalactoferrin, Lactoferricin-H, Lfcin-H, Kaliocin-1, Lactoferroxin-A, Lactoferroxin-B, Lactoferroxin-C, LTF, GIG12, LF

Target/Specificity

Human LTF / Lactoferrin.

Reconstitution & Storage

Store at -20°C. Minimize freezing and thawing.

Precautions

LF / LTF / Lactoferrin Antibody (Internal) is for research use only and not for use in diagnostic or therapeutic procedures.

LF / LTF / Lactoferrin Antibody (Internal) - Protein Information

Name LTF (HGNC:6720)

Synonyms GIG12, LF

Function

Transferrins are iron binding transport proteins which can bind two Fe(3+) ions in association with the binding of an anion, usually bicarbonate.

Cellular Location

[Isoform 1]: Secreted. Cytoplasmic granule. Note=Secreted into most exocrine fluids by various endothelial cells Stored in the secondary granules of neutrophils

Tissue Location

High levels are found in saliva and tears, intermediate levels in serum and plasma, and low levels in urine. In kidney, detected in the distal collecting tubules in the medulla but not in the cortical




region or in blood vessels. Detected in peripheral blood neutrophils (at protein level). Isoform 1 and isoform DeltaLf are expressed in breast, prostate, spleen, pancreas, kidney, small intestine, lung, skeletal muscle, uterus, thymus and fetal liver Isoform 1 is expressed in brain, testis and peripheral blood leukocytes; isoform DeltaLf is barely detectable in these tissues Isoform DeltaLf is expressed in placenta, liver and ovary; isoform 1 is barely detectable in these tissues. In kidney, isoform 1 is expressed at high levels in the collecting tubules of the medulla but at very low levels in the cortex.

LF / LTF / Lactoferrin Antibody (Internal) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

LF / LTF / Lactoferrin Antibody (Internal) - Images



Antibody (0.05 ug/ml) staining of Peripheral Blood Mononucleocyte lysate (35 ug protein in RIPA...



Anti-LTF / Lactoferrin antibody IHC of human neutrophils.

LF / LTF / Lactoferrin Antibody (Internal) - Background





Transferrins are iron binding transport proteins which can bind two Fe(3+) ions in association with the binding of an anion, usually bicarbonate. Lactoferricin binds to the bacterial surface and is crucial for the bactericidal functions. Has some antiviral activity against papillomavirus infection. N-terminal region shows strong antifungal activity against C.albicans. Contains two BBXB heparin-binding consensus sequences that appear to form the predominate functional GAG-binding site. Lactoferroxins A, B and C have opioid antagonist activity. Lactoferroxin A shows preference for mu-receptors, while lactoferroxin B and C have somewhat higher degrees of preference for kappa-receptors than for mu-receptors. Isoform DeltaLf: transcription factor with antiproliferative properties and ability to induce cell cycle arrest. Binds to the DeltaLf response element found in the SKP1, BAX, DCPS, and SELH promoters.

LF / LTF / Lactoferrin Antibody (Internal) - References

Rey M.W.,et al.Nucleic Acids Res. 18:5288-5288(1990).
Cho Y.Y.,et al.Thesis (1994), Genetic Engineering Research Institute, South Korea.
Siebert P.D.,et al.Proc. Natl. Acad. Sci. U.S.A. 94:2198-2203(1997).
Cheng H.,et al.Sheng Wu Gong Cheng Xue Bao 17:385-387(2001).
Kumar J.,et al.Indian J. Biochem. Biophys. 40:14-21(2003).