

Myelin Basic Protein / MBP Antibody (N-Terminus, clone V/h5)
Mouse Monoclonal Antibody
Catalog # ALS12758**Specification**

Myelin Basic Protein / MBP Antibody (N-Terminus, clone V/h5) - Product Information

Application	IHC
Primary Accession	P02686
Reactivity	Human, Rabbit, Sheep, Bovine, Guinea Pig
Host	Mouse
Clonality	Monoclonal
Calculated MW	33kDa KDa

Myelin Basic Protein / MBP Antibody (N-Terminus, clone V/h5) - Additional Information**Gene ID** 4155**Other Names**

Myelin basic protein, MBP, Myelin A1 protein, Myelin membrane encephalitogenic protein, MBP

Target/Specificity

Recognizes predominantly residues 45-91

Reconstitution & Storage

+4°C or -20°C, Avoid repeated freezing and thawing.

Precautions

Myelin Basic Protein / MBP Antibody (N-Terminus, clone V/h5) is for research use only and not for use in diagnostic or therapeutic procedures.

Myelin Basic Protein / MBP Antibody (N-Terminus, clone V/h5) - Protein Information**Name** MBP**Function**

The classic group of MBP isoforms (isoform 4-isoform 14) are with PLP the most abundant protein components of the myelin membrane in the CNS. They have a role in both its formation and stabilization. The smaller isoforms might have an important role in remyelination of denuded axons in multiple sclerosis. The non-classic group of MBP isoforms (isoform 1-isoform 3/Golli-MBPs) may preferentially have a role in the early developing brain long before myelination, maybe as components of transcriptional complexes, and may also be involved in signaling pathways in T-cells and neural cells. Differential splicing events combined with optional post-translational modifications give a wide spectrum of isomers, with each of them potentially having a specialized function. Induces T-cell proliferation.

Cellular Location

Myelin membrane; Peripheral membrane protein; Cytoplasmic side. Note=Cytoplasmic side of myelin


Tissue Location

MBP isoforms are found in both the central and the peripheral nervous system, whereas Golli-MBP isoforms are expressed in fetal thymus, spleen and spinal cord, as well as in cell lines derived from the immune system.

Myelin Basic Protein / MBP Antibody (N-Terminus, clone V/h5) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Myelin Basic Protein / MBP Antibody (N-Terminus, clone V/h5) - Images

Anti-Myelin Basic Protein antibody IHC of human brain, cerebellum.

Myelin Basic Protein / MBP Antibody (N-Terminus, clone V/h5) - Background

The classic group of MBP isoforms (isoform 4-isoform 14) are with PLP the most abundant protein components of the myelin membrane in the CNS. They have a role in both its formation and stabilization. The smaller isoforms might have an important role in remyelination of denuded axons in multiple sclerosis. The non- classic group of MBP isoforms (isoform 1-isoform 3/Golli-MBPs) may preferentially have a role in the early developing brain long before myelination, maybe as components of transcriptional complexes, and may also be involved in signaling pathways in T-cells and neural cells. Differential splicing events combined with optional post-translational modifications give a wide spectrum of isomers, with each of them potentially having a specialized function. Induces T-cell proliferation.

Myelin Basic Protein / MBP Antibody (N-Terminus, clone V/h5) - References

- Carnegie P.R., et al. Biochem. J. 123:57-67(1971).
Roth H.J., et al. J. Neurosci. Res. 16:227-238(1986).
Kamholz J., et al. Proc. Natl. Acad. Sci. U.S.A. 83:4962-4966(1986).
Roth H.J., et al. J. Neurosci. Res. 17:321-328(1987).

Streicher R., et al. Biol. Chem. Hoppe-Seyler 370:503-510(1989).