

**HSPA5 / GRP78 / BiP Antibody (aa525-628, clone 4E3)**  
**Mouse Monoclonal Antibody**  
**Catalog # ALS13203****Specification****HSPA5 / GRP78 / BiP Antibody (aa525-628, clone 4E3) - Product Information**

Application	IHC
Primary Accession	<a href="#">P11021</a>
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Calculated MW	72kDa KDa

**HSPA5 / GRP78 / BiP Antibody (aa525-628, clone 4E3) - Additional Information****Gene ID** 3309**Other Names**

78 kDa glucose-regulated protein, GRP-78, Endoplasmic reticulum luminal Ca(2+)-binding protein grp78, Heat shock 70 kDa protein 5, Immunoglobulin heavy chain-binding protein, BiP, HSPA5, GRP78

**Target/Specificity**

Human HSPA5

**Reconstitution & Storage**

Long term: -20°C; Short term: +4°C. Avoid repeat freeze-thaw cycles.

**Precautions**

HSPA5 / GRP78 / BiP Antibody (aa525-628, clone 4E3) is for research use only and not for use in diagnostic or therapeutic procedures.

**HSPA5 / GRP78 / BiP Antibody (aa525-628, clone 4E3) - Protein Information****Name** HSPA5 ([HGNC:5238](#))**Function**

Endoplasmic reticulum chaperone that plays a key role in protein folding and quality control in the endoplasmic reticulum lumen (PubMed:<a href="http://www.uniprot.org/citations/2294010" target="\_blank">2294010</a>, PubMed:<a href="http://www.uniprot.org/citations/23769672" target="\_blank">23769672</a>, PubMed:<a href="http://www.uniprot.org/citations/23990668" target="\_blank">23990668</a>, PubMed:<a href="http://www.uniprot.org/citations/28332555" target="\_blank">28332555</a>). Involved in the correct folding of proteins and degradation of misfolded proteins via its interaction with DNAJC10/ERdj5, probably to facilitate the release of DNAJC10/ERdj5 from its substrate (By similarity). Acts as a key repressor of the ERN1/IRE1-mediated unfolded protein response (UPR) (PubMed:<a href="http://www.uniprot.org/citations/1550958" target="\_blank">1550958</a>, PubMed:<a href="http://www.uniprot.org/citations/19538957" target="\_blank">19538957</a>). In the

unstressed endoplasmic reticulum, recruited by DNAJB9/ERdj4 to the luminal region of ERN1/IRE1, leading to disrupt the dimerization of ERN1/IRE1, thereby inactivating ERN1/IRE1 (By similarity). Accumulation of misfolded protein in the endoplasmic reticulum causes release of HSPA5/BiP from ERN1/IRE1, allowing homodimerization and subsequent activation of ERN1/IRE1 (By similarity). Plays an auxiliary role in post-translational transport of small presecretory proteins across endoplasmic reticulum (ER). May function as an allosteric modulator for SEC61 channel-forming translocon complex, likely cooperating with SEC62 to enable the productive insertion of these precursors into SEC61 channel. Appears to specifically regulate translocation of precursors having inhibitory residues in their mature region that weaken channel gating. May also play a role in apoptosis and cell proliferation (PubMed:<a href="http://www.uniprot.org/citations/26045166" target="\_blank">26045166</a>).

#### **Cellular Location**

Endoplasmic reticulum lumen. Melanosome. Cytoplasm {ECO:0000250|UniProtKB:P20029}. Cell surface Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV (PubMed:12643545). Localizes to the cell surface of epithelial cells in response to high levels of free iron (PubMed:20484814, PubMed:24355926, PubMed:27159390)

#### **Volume**


50 µl

#### **HSPA5 / GRP78 / BiP Antibody (aa525-628, clone 4E3) - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **HSPA5 / GRP78 / BiP Antibody (aa525-628, clone 4E3) - Images**



Anti-HSPA5 / GRP78 antibody IHC of human breast.



Anti-HSPA5 / GRP78 antibody IHC of human thyroid.

#### **HSPA5 / GRP78 / BiP Antibody (aa525-628, clone 4E3) - Background**

Probably plays a role in facilitating the assembly of multimeric protein complexes inside the endoplasmic reticulum. Involved in the correct folding of proteins and degradation of misfolded proteins via its interaction with DNAJC10, probably to facilitate the release of DNAJC10 from its substrate.

#### **HSPA5 / GRP78 / BiP Antibody (aa525-628, clone 4E3) - References**

Ting J.,et al.DNA 7:275-286(1988).  
Chao C.C.K.,et al.Submitted (DEC-1995) to the EMBL/GenBank/DDBJ databases.  
Hansen J.J.,et al.Submitted (JAN-2000) to the EMBL/GenBank/DDBJ databases.  
Bermudez-Fajardo A.,et al.Submitted (DEC-1999) to the EMBL/GenBank/DDBJ databases.  
Humphray S.J.,et al.Nature 429:369-374(2004).