

**EMA / MUC1 Antibody**  
**Rabbit Polyclonal Antibody**  
**Catalog # ALS13269****Specification**

---

**EMA / MUC1 Antibody - Product Information**

Application	IHC-P
Primary Accession	<a href="#">P15941</a>
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	122kDa KDa
Dilution	IHC-P~~N/A

**EMA / MUC1 Antibody - Additional Information****Gene ID** 4582**Other Names**

Mucin-1, MUC-1, Breast carcinoma-associated antigen DF3, Cancer antigen 15-3, CA 15-3, Carcinoma-associated mucin, Episialin, H23AG, Krebs von den Lungen-6, KL-6, PEMT, Peanut-reactive urinary mucin, PUM, Polymorphic epithelial mucin, PEM, Tumor-associated epithelial membrane antigen, EMA, Tumor-associated mucin, CD227, Mucin-1 subunit alpha, MUC1-NT, MUC1-alpha, Mucin-1 subunit beta, MUC1-beta, MUC1-CT, MUC1, PUM

**Reconstitution & Storage**

Long term: -20°C; Short term: +4°C. Avoid repeat freeze-thaw cycles.

**Precautions**

EMA / MUC1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**EMA / MUC1 Antibody - Protein Information****Name** MUC1**Synonyms** PUM**Function**

The alpha subunit has cell adhesive properties. Can act both as an adhesion and an anti-adhesion protein. May provide a protective layer on epithelial cells against bacterial and enzyme attack.

**Cellular Location**

Apical cell membrane; Single-pass type I membrane protein. Note=Exclusively located in the apical domain of the plasma membrane of highly polarized epithelial cells After endocytosis, internalized and recycled to the cell membrane Located to microvilli and to the tips of long filopodial protusions [Isoform Y]: Secreted. [Mucin-1 subunit beta]: Cell membrane. Cytoplasm. Nucleus. Note=On EGF and PDGFRB stimulation, transported to the nucleus through interaction

with CTNNB1, a process which is stimulated by phosphorylation. On HRG stimulation, colocalizes with JUP/gamma-catenin at the nucleus

**Tissue Location**

Expressed on the apical surface of epithelial cells, especially of airway passages, breast and uterus. Also expressed in activated and unactivated T-cells. Overexpressed in epithelial tumors, such as breast or ovarian cancer and also in non-epithelial tumor cells. Isoform Y is expressed in tumor cells only


**Volume**

500 µl


**EMA / MUC1 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**EMA / MUC1 Antibody - Images**

Anti-MUC1 antibody IHC of human kidney.



Anti-MUC1 antibody IHC of human pancreas.

#### **EMA / MUC1 Antibody - Background**

The alpha subunit has cell adhesive properties. Can act both as an adhesion and an anti-adhesion protein. May provide a protective layer on epithelial cells against bacterial and enzyme attack.

#### **EMA / MUC1 Antibody - References**

- Lan M.S.,et al.J. Biol. Chem. 265:15294-15299(1990).
- Ligtenberg M.J.L.,et al.J. Biol. Chem. 265:5573-5578(1990).
- Gendler S.J.,et al.J. Biol. Chem. 265:15286-15293(1990).
- Lancaster C.A.,et al.Biochem. Biophys. Res. Commun. 173:1019-1029(1990).
- Wreschner D.H.,et al.Eur. J. Biochem. 189:463-473(1990).