

### HMG2 / HMGB2 Antibody (clone 3C7)

Mouse Monoclonal Antibody Catalog # ALS13339

#### **Specification**

### HMG2 / HMGB2 Antibody (clone 3C7) - Product Information

WB, IHC-P, E Application **Primary Accession** P26583 Reactivity Human Host Mouse Clonality **Monoclonal** Calculated MW 24kDa KDa Dilution WB~~1:1000 IHC-P~~N/A E~~N/A

#### HMG2 / HMGB2 Antibody (clone 3C7) - Additional Information

**Gene ID 3148** 

# **Other Names**

High mobility group protein B2, High mobility group protein 2, HMG-2, HMGB2, HMG2

#### **Reconstitution & Storage**

Store at -20°C. Aliquot to avoid freeze/thaw cycles.

#### **Precautions**

HMG2 / HMGB2 Antibody (clone 3C7) is for research use only and not for use in diagnostic or therapeutic procedures.

# HMG2 / HMGB2 Antibody (clone 3C7) - Protein Information

#### Name HMGB2

### Synonyms HMG2

#### **Function**

Multifunctional protein with various roles in different cellular compartments. May act in a redox sensitive manner. In the nucleus is an abundant chromatin-associated non-histone protein involved in transcription, chromatin remodeling and V(D)J recombination and probably other processes. Binds DNA with a preference to non- canonical DNA structures such as single-stranded DNA. Can bent DNA and enhance DNA flexibility by looping thus providing a mechanism to promote activities on various gene promoters by enhancing transcription factor binding and/or bringing distant regulatory sequences into close proximity (PubMed:<a

href="http://www.uniprot.org/citations/11909973" target="\_blank">11909973</a>, PubMed:<a href="http://www.uniprot.org/citations/18413230" target="\_blank">18413230</a>, PubMed:<a href="http://www.uniprot.org/citations/19522541" target="\_blank">19522541</a>, PubMed:<a href="http://www.uniprot.org/citations/19965638" target="\_blank">19965638</a>, PubMed:<a



href="http://www.uniprot.org/citations/20123072" target="\_blank">20123072</a>, PubMed:<a href="http://www.uniprot.org/citations/7797075" target="\_blank">7797075</a>). Involved in V(D)J recombination by acting as a cofactor of the RAG complex: acts by stimulating cleavage and RAG protein binding at the 23 bp spacer of conserved recombination signal sequences (RSS) (By similarity). Proposed to be involved in the innate immune response to nucleic acids by acting as a promiscuous immunogenic DNA/RNA sensor which cooperates with subsequent discriminative sensing by specific pattern recognition receptors (By similarity). In the extracellular compartment acts as a chemokine. Promotes proliferation and migration of endothelial cells implicating AGER/RAGE (PubMed:<a href="http://www.uniprot.org/citations/19811285" target="\_blank">19811285</a>). Has antimicrobial activity in gastrointestinal epithelial tissues (PubMed:<a href="http://www.uniprot.org/citations/23877675" target="\_blank">23877675</a>). Involved in inflammatory response to antigenic stimulus coupled with pro- inflammatory activity (By similarity). Involved in modulation of neurogenesis probably by regulation of neural stem proliferation (By similarity). Involved in articular cartilage surface maintenance implicating LEF1 and the Wnt/beta-catenin pathway (By similarity).

#### **Cellular Location**

Nucleus. Chromosome. Cytoplasm. Secreted. Note=In basal state predominantly nuclear.

#### **Tissue Location**


Expressed in gastric and intestinal tissues (at protein level).

# HMG2 / HMGB2 Antibody (clone 3C7) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

#### HMG2 / HMGB2 Antibody (clone 3C7) - Images



HMGB2 monoclonal antibody (M03), clone 3C7 Western blot of HMGB2 expression in HeLa NE.





Western blot of HMGB2 expression in transfected 293T cell line by HMGB2 monoclonal antibody...

# HMG2 / HMGB2 Antibody (clone 3C7) - Background

DNA binding proteins that associates with chromatin and has the ability to bend DNA. Binds preferentially single-stranded DNA. Involved in V(D)J recombination by acting as a cofactor of the RAG complex. Acts by stimulating cleavage and RAG protein binding at the 23 bp spacer of conserved recombination signal sequences (RSS) (By similarity).

# HMG2 / HMGB2 Antibody (clone 3C7) - References

Majumdar A., et al. Nucleic Acids Res. 19:6643-6643(1991). Shirakawa H., et al.J. Biol. Chem. 267:6641-6645(1992). Kalnine N., et al. Submitted (OCT-2004) to the EMBL/GenBank/DDBJ databases. Ota T., et al. Nat. Genet. 36:40-45(2004). Mural R.J., et al. Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.