

**FNIP2 Antibody (C-Terminus)**  
**Rabbit Polyclonal Antibody**  
**Catalog # ALS13522****Specification****FNIP2 Antibody (C-Terminus) - Product Information**

Application	WB, IHC
Primary Accession	<a href="#">Q9P278</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	122kDa KDa

**FNIP2 Antibody (C-Terminus) - Additional Information****Gene ID** 57600**Other Names**

Folliculin-interacting protein 2, FNIP1-like protein, O6-methylguanine-induced apoptosis 1 protein, FNIP2, FNIPL, KIAA1450, MAPO1

**Target/Specificity**

Human FNIP2

**Reconstitution & Storage**

Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles. Store undiluted.

**Precautions**

FNIP2 Antibody (C-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

**FNIP2 Antibody (C-Terminus) - Protein Information****Name** FNIP2 {ECO:0000303|PubMed:18663353, ECO:0000312|HGNC:HGNC:29280}**Function**

Binding partner of the GTPase-activating protein FLCN: involved in the cellular response to amino acid availability by regulating the non-canonical mTORC1 signaling cascade controlling the MiT/TFE factors TFEB and TFE3 (PubMed:<a href="http://www.uniprot.org/citations/18663353" target="\_blank">18663353</a>, PubMed:<a href="http://www.uniprot.org/citations/31672913" target="\_blank">31672913</a>, PubMed:<a href="http://www.uniprot.org/citations/36103527" target="\_blank">36103527</a>). Required to promote FLCN recruitment to lysosomes and interaction with Rag GTPases, leading to activation of the non- canonical mTORC1 signaling (By similarity). In low-amino acid conditions, component of the lysosomal folliculin complex (LFC) on the membrane of lysosomes, which inhibits the GTPase-activating activity of FLCN, thereby inactivating mTORC1 and promoting nuclear translocation of TFEB and TFE3 (PubMed:<a href="http://www.uniprot.org/citations/31672913" target="\_blank">31672913</a>, PubMed:<a href="http://www.uniprot.org/citations/36103527" target="\_blank">36103527</a>). Upon amino

acid restimulation, disassembly of the LFC complex liberates the GTPase- activating activity of FLCN, leading to activation of mTORC1 and subsequent inactivation of TFEB and TFE3 (PubMed:<a href="http://www.uniprot.org/citations/31672913" target="\_blank">31672913</a>). Together with FLCN, regulates autophagy: following phosphorylation by ULK1, interacts with GABARAP and promotes autophagy (PubMed:<a href="http://www.uniprot.org/citations/25126726" target="\_blank">25126726</a>). In addition to its role in mTORC1 signaling, also acts as a co-chaperone of HSP90AA1/Hsp90: inhibits the ATPase activity of HSP90AA1/Hsp90, leading to activate both kinase and non-kinase client proteins of HSP90AA1/Hsp90 (PubMed:<a href="http://www.uniprot.org/citations/18403135" target="\_blank">18403135</a>). Acts as a scaffold to load client protein FLCN onto HSP90AA1/Hsp90 (PubMed:<a href="http://www.uniprot.org/citations/18403135" target="\_blank">18403135</a>). Competes with the activating co-chaperone AHSA1 for binding to HSP90AA1, thereby providing a reciprocal regulatory mechanism for chaperoning of client proteins (PubMed:<a href="http://www.uniprot.org/citations/18403135" target="\_blank">18403135</a>). May play a role in the signal transduction pathway of apoptosis induced by O6-methylguanine-mispaired lesions (By similarity).

### Cellular Location

Lysosome membrane. Cytoplasm. Note=Colocalizes with FLCN in the cytoplasm.

### Tissue Location

Widely expressed with highest levels in muscle, nasal mucosa, salivary gland, uvula, fat, liver, heart, placenta and pancreas (PubMed:18403135, PubMed:18663353, PubMed:27353360) Moderately expressed in the lung, small intestine, kidney and brain Lower levels detected in renal cell carcinoma than in normal kidney tissue (PubMed:18403135). Higher levels detected in oncocytoma tumors than in normal kidney. Higher levels detected in renal cell carcinoma tumors than in normal kidney tissue (PubMed:27353360)


### FNIP2 Antibody (C-Terminus) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### FNIP2 Antibody (C-Terminus) - Images






Western blot of FNIP2 in rat skeletal muscle lysate with FNIP2 antibody at (A) 1 and (B) 2 ug/ml.



Anti-FNIP2 antibody IHC of human colon, submucosal plexus.



Anti-FNIP2 antibody IHC of human placenta.

#### **FNIP2 Antibody (C-Terminus) - Background**

May play a role in the signal transduction pathway of apoptosis induced by O6-methylguanine-mispaired lesions (By similarity). May be involved in energy and/or nutrient sensing through the AMPK and mTOR signaling pathways. May regulate phosphorylation of RPS6KB1.

#### **FNIP2 Antibody (C-Terminus) - References**

Nagase T., et al. DNA Res. 7:143-150(2000).

Ota T.,et al.Nat. Genet. 36:40-45(2004).  
Hasumi H.,et al.Gene 415:60-67(2008).  
Takagi Y.,et al.Oncogene 27:5339-5347(2008).  
Komori K.,et al.Oncogene 28:1142-1150(2009).