

ZNF521 Antibody (C-Terminus)
Rabbit Polyclonal Antibody
Catalog # ALS13591**Specification**

ZNF521 Antibody (C-Terminus) - Product Information

Application	IHC-P
Primary Accession	Q96K83
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	148kDa KDa
Dilution	IHC-P~~N/A

ZNF521 Antibody (C-Terminus) - Additional Information**Gene ID** 25925**Other Names**

Zinc finger protein 521, Early hematopoietic zinc finger protein, LYST-interacting protein 3, ZNF521, EHZF, LIP3

Reconstitution & Storage

Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles. Store undiluted.

Precautions

ZNF521 Antibody (C-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

ZNF521 Antibody (C-Terminus) - Protein Information**Name** ZNF521**Synonyms** EHZF, LIP3**Function**

Transcription factor that can both act as an activator or a repressor depending on the context. Involved in BMP signaling and in the regulation of the immature compartment of the hematopoietic system. Associates with SMADs in response to BMP2 leading to activate transcription of BMP target genes. Acts as a transcriptional repressor via its interaction with EBF1, a transcription factor involved specification of B-cell lineage; this interaction preventing EBF1 to bind DNA and activate target genes.

Cellular Location

Nucleus.

Tissue Location

Predominantly expressed in hematopoietic cells. Present in organs and tissues that contain stem


and progenitor cells, myeloid and/or lymphoid: placenta, spleen, lymph nodes, thymus, bone marrow and fetal liver. Within the hematopoietic system, it is abundant in CD34(+) cells but undetectable in mature peripheral blood leukocytes, and its levels rapidly decrease during the differentiation of CD34(+) cells in response to hemopoietins

ZNF521 Antibody (C-Terminus) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ZNF521 Antibody (C-Terminus) - Images



Anti-ZNF521 antibody IHC of human small intestine.



Anti-ZNF521 antibody IHC of human kidney.

ZNF521 Antibody (C-Terminus) - Background

Transcription factor that can both act as an activator or a repressor depending on the context.

Involved in BMP signaling and in the regulation of the immature compartment of the hematopoietic system. Associates with SMADs in response to BMP2 leading to activate transcription of BMP target genes. Acts as a transcriptional repressor via its interaction with EBF1, a transcription factor involved specification of B-cell lineage; this interaction preventing EBF1 to bind DNA and activate target genes.

ZNF521 Antibody (C-Terminus) - References

Bond H.M.,et al.Blood 103:2062-2070(2004).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.
Mullighan C.G.,et al.Nature 446:758-764(2007).
Bechtel S.,et al.BMC Genomics 8:399-399(2007).