

Basigin / Emmprin / CD147 Antibody
Rat Monoclonal Antibody
Catalog # ALS13674**Specification**

Basigin / Emmprin / CD147 Antibody - Product Information

Application	IHC
Primary Accession	P35613
Reactivity	Mouse
Host	Rat
Clonality	Monoclonal
Calculated MW	42kDa KDa

Basigin / Emmprin / CD147 Antibody - Additional Information**Gene ID** 682**Other Names**

Basigin, 5F7, Collagenase stimulatory factor, Extracellular matrix metalloproteinase inducer, EMMPRIN, Leukocyte activation antigen M6, OK blood group antigen, Tumor cell-derived collagenase stimulatory factor, TCSF, CD147, BSG

Target/Specificity

Recognizes mouse CD147 (also known as basigin). Clone recognizes both isoforms of CD147, the common form containing 2 Ig-like domains and the rare form containing 3 Ig-like domains.

Reconstitution & Storage

Long term: -20°C; Short term: +4°C; Avoid freeze-thaw cycles.

Precautions

Basigin / Emmprin / CD147 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Basigin / Emmprin / CD147 Antibody - Protein Information**Name** BSG ([HGNC:1116](#))**Function**

[Isoform 1]: Essential for normal retinal maturation and development (By similarity). Acts as a retinal cell surface receptor for NXNL1 and plays an important role in NXNL1-mediated survival of retinal cone photoreceptors (PubMed:25957687). In association with glucose transporter SLC16A1/GLUT1 and NXNL1, promotes retinal cone survival by enhancing aerobic glycolysis and accelerating the entry of glucose into photoreceptors (PubMed:25957687). May act as a potent stimulator of IL6 secretion in multiple cell lines that include monocytes (PubMed:21620857).

Cellular Location

Melanosome. Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV. [Isoform 2]: Cell membrane; Single-pass type I membrane protein {ECO:0000250|UniProtKB:P26453}. Endosome Endoplasmic reticulum membrane; Single-pass type I membrane protein {ECO:0000250|UniProtKB:P26453} Basolateral cell membrane; Single-pass type I membrane protein {ECO:0000250|UniProtKB:P26453} [Isoform 4]: Cell membrane; Single-pass type I membrane protein {ECO:0000250|UniProtKB:P26453}

Tissue Location

[Isoform 1]: Retina-specific (PubMed:25957687). Expressed in retinal cone photoreceptors (at protein level) (PubMed:25957687). [Isoform 3]: Highly expressed in the bone marrow, fetal liver, lung, testis and thymus.


Volume

50 µl

Basigin / Emmprin / CD147 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Basigin / Emmprin / CD147 Antibody - Images

Anti-CD147 antibody IHC of mouse kidney.

Basigin / Emmprin / CD147 Antibody - Background

Plays an important role in targeting the monocarboxylate transporters SLC16A1, SLC16A3 and SLC16A8 to the plasma membrane. Plays pivotal roles in spermatogenesis, embryo implantation, neural network formation and tumor progression. Stimulates adjacent fibroblasts to produce matrix metalloproteinases (MMPs). Seems to be a receptor for oligomannosidic glycans. In vitro, promotes outgrowth of astrocytic processes.

Basigin / Emmprin / CD147 Antibody - References

Kasinrerk W.,et al.J. Immunol. 149:847-854(1992).
Miyauchi T.,et al.J. Biochem. 110:770-774(1991).
Biswas C.,et al.Cancer Res. 55:434-439(1995).
Wakasugi H.,et al.Submitted (MAR-1996) to the EMBL/GenBank/DDBJ databases.
Decastro R.,et al.Submitted (MAR-1996) to the EMBL/GenBank/DDBJ databases.