

**STEAP4 Antibody (aa201-250)**  
**Rabbit Polyclonal Antibody**  
**Catalog # ALS13679****Specification****STEAP4 Antibody (aa201-250) - Product Information**

Application	IHC-P, IF, E
Primary Accession	<a href="#">Q687X5</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	52kDa KDa
Dilution	IHC-P~~N/A IF~~1:50~200 E~~N/A

**STEAP4 Antibody (aa201-250) - Additional Information****Gene ID 79689****Other Names**

Metalloreductase STEAP4, 1.16.1.-, Six-transmembrane epithelial antigen of prostate 4, SixTransMembrane protein of prostate 2, Tumor necrosis factor, alpha-induced protein 9, STEAP4, STAMP2, TNFAIP9

**Target/Specificity**

STEAP4 Antibody detects endogenous levels of total STEAP4 protein.

**Reconstitution & Storage**

Store at -20°C for up to one year.

**Precautions**

STEAP4 Antibody (aa201-250) is for research use only and not for use in diagnostic or therapeutic procedures.

**STEAP4 Antibody (aa201-250) - Protein Information****Name STEAP4****Synonyms** STAMP2 {ECO:0000303|PubMed:15897894}, TN**Function**

Integral membrane protein that functions as a NADPH-dependent ferric-chelate reductase, using NADPH from one side of the membrane to reduce a Fe(3+) chelate that is bound on the other side of the membrane. Mediates sequential transmembrane electron transfer from NADPH to FAD and onto heme, and finally to the Fe(3+) chelate (PubMed:<a href="http://www.uniprot.org/citations/30337524" target="\_blank">30337524</a>). Can also reduce Cu(2+) to Cu(1+) (By similarity). Plays a role in systemic metabolic homeostasis,

integrating inflammatory and metabolic responses (By similarity). Associated with obesity and insulin-resistance (PubMed:<a href="http://www.uniprot.org/citations/18381574" target="\_blank">18381574</a>, PubMed:<a href="http://www.uniprot.org/citations/18430367" target="\_blank">18430367</a>). Involved in inflammatory arthritis, through the regulation of inflammatory cytokines (PubMed:<a href="http://www.uniprot.org/citations/19660107" target="\_blank">19660107</a>). Inhibits anchorage-independent cell proliferation (PubMed:<a href="http://www.uniprot.org/citations/19787193" target="\_blank">19787193</a>).

**Cellular Location**

Cell membrane; Multi-pass membrane protein. Golgi apparatus membrane; Multi-pass membrane protein. Early endosome membrane; Multi-pass membrane protein

**Tissue Location**

Ubiquitous. Highly expressed in adipose tissue. Expressed in placenta, lung, heart and prostate. Detected at lower levels in liver, skeletal muscle, pancreas, testis and small intestine. Highly expressed in joints of patients with rheumatoid arthritis and localized with CD68 cells, a marker for macrophages


**Volume**

50 µl


**STEAP4 Antibody (aa201-250) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**STEAP4 Antibody (aa201-250) - Images**

Immunofluorescence of HeLa cells, using STEAP4 Antibody.



Anti-STEAP4 antibody IHC of human liver.

#### **STEAP4 Antibody (aa201-250) - Background**

Metalloreductase that has the ability to reduce both Fe(3+) to Fe(2+) and Cu(2+) to Cu(1+). Uses NAD(+) as acceptor. Plays a role in systemic metabolic homeostasis, integrating inflammatory and metabolic responses (By similarity). Associated with obesity and insulin-resistance. Involved in inflammatory arthritis, through the regulation of inflammatory cytokines. Inhibits anchorage-independent cell proliferation.

#### **STEAP4 Antibody (aa201-250) - References**

- Korkmaz C.G., et al. Oncogene 24:4934-4945(2005).
- Ota T., et al. Nat. Genet. 36:40-45(2004).
- Bechtel S., et al. BMC Genomics 8:399-399(2007).
- Ohgami R.S., et al. Nat. Genet. 37:1264-1269(2005).
- Zhang C.M., et al. Acta Pharmacol. Sin. 29:587-592(2008).