

**SATB2 Antibody (C-Terminus)**  
**Rabbit Polyclonal Antibody**  
**Catalog # ALS13726****Specification**

---

**SATB2 Antibody (C-Terminus) - Product Information**

Application	IF, IHC
Primary Accession	<a href="#">Q9UPW6</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	83kDa KDa

**SATB2 Antibody (C-Terminus) - Additional Information****Gene ID** 23314**Other Names**

DNA-binding protein SATB2, Special AT-rich sequence-binding protein 2, SATB2, KIAA1034

**Reconstitution & Storage**

Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles. Store undiluted.

**Precautions**

SATB2 Antibody (C-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

**SATB2 Antibody (C-Terminus) - Protein Information****Name** SATB2**Synonyms** KIAA1034**Function**

Binds to DNA, at nuclear matrix- or scaffold-associated regions. Thought to recognize the sugar-phosphate structure of double- stranded DNA. Transcription factor controlling nuclear gene expression, by binding to matrix attachment regions (MARs) of DNA and inducing a local chromatin-loop remodeling. Acts as a docking site for several chromatin remodeling enzymes and also by recruiting corepressors (HDACs) or coactivators (HATs) directly to promoters and enhancers. Required for the initiation of the upper-layer neurons (UL1) specific genetic program and for the inactivation of deep-layer neurons (DL) and UL2 specific genes, probably by modulating BCL11B expression. Repressor of Ctip2 and regulatory determinant of corticocortical connections in the developing cerebral cortex. May play an important role in palate formation. Acts as a molecular node in a transcriptional network regulating skeletal development and osteoblast differentiation.

**Cellular Location**

Nucleus matrix {ECO:0000255|PROSITE- ProRule:PRU00108,

ECO:0000255|PROSITE-ProRule:PRU00374, ECO:0000269|PubMed:14701874}

#### **Tissue Location**


High expression in adult brain, moderate expression in fetal brain, and weak expression in adult liver, kidney, and spinal cord and in select brain regions, including amygdala, corpus callosum, caudate nucleus, and hippocampus.

#### **SATB2 Antibody (C-Terminus) - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **SATB2 Antibody (C-Terminus) - Images**



Immunofluorescence of SATB2 in Human Brain cells with SATB2 antibody at 50 ug/ml.



Anti-SATB2 antibody IHC of human brain, cortex.

**SATB2 Antibody (C-Terminus) - Background**

Binds to DNA, at nuclear matrix- or scaffold-associated regions. Thought to recognize the sugar-phosphate structure of double-stranded DNA. Transcription factor controlling nuclear gene expression, by binding to matrix attachment regions (MARs) of DNA and inducing a local chromatin-loop remodeling. Acts as a docking site for several chromatin remodeling enzymes and also by recruiting corepressors (HDACs) or coactivators (HATs) directly to promoters and enhancers. Required for the initiation of the upper- layer neurons (UL1) specific genetic program and for the inactivation of deep-layer neurons (DL) and UL2 specific genes, probably by modulating BCL11B expression. Repressor of Ctip2 and regulatory determinant of corticocortical connections in the developing cerebral cortex. May play an important role in palate formation. Acts as a molecular node in a transcriptional network regulating skeletal development and osteoblast differentiation.

**SATB2 Antibody (C-Terminus) - References**

Kikuno R.,et al.DNA Res. 6:197-205(1999).  
Ota T.,et al.Nat. Genet. 36:40-45(2004).  
Hillier L.W.,et al.Nature 434:724-731(2005).  
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.  
Dobrev G.,et al.Genes Dev. 17:3048-3061(2003).