

**CDKN1B / p27 Kip1 Antibody (C-Terminus)**  
**Rabbit Polyclonal Antibody**  
**Catalog # ALS13909****Specification**

---

**CDKN1B / p27 Kip1 Antibody (C-Terminus) - Product Information**

Application	IHC
Primary Accession	<a href="#">P46527</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	22kDa KDa

**CDKN1B / p27 Kip1 Antibody (C-Terminus) - Additional Information****Gene ID** 1027**Other Names**

Cyclin-dependent kinase inhibitor 1B, Cyclin-dependent kinase inhibitor p27, p27Kip1, CDKN1B, KIP1

**Reconstitution & Storage**

Store at 2°C to 8°C degrees. Do not freeze.

**Precautions**

CDKN1B / p27 Kip1 Antibody (C-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

**CDKN1B / p27 Kip1 Antibody (C-Terminus) - Protein Information****Name** CDKN1B {ECO:0000303|PubMed:20824794}**Function**

Important regulator of cell cycle progression. Inhibits the kinase activity of CDK2 bound to cyclin A, but has little inhibitory activity on CDK2 bound to SPDYA (PubMed:<a href="http://www.uniprot.org/citations/28666995" target="\_blank">28666995</a>). Involved in G1 arrest. Potent inhibitor of cyclin E- and cyclin A-CDK2 complexes. Forms a complex with cyclin type D-CDK4 complexes and is involved in the assembly, stability, and modulation of CCND1-CDK4 complex activation. Acts either as an inhibitor or an activator of cyclin type D-CDK4 complexes depending on its phosphorylation state and/or stoichiometry.

**Cellular Location**

Nucleus. Cytoplasm. Endosome. Note=Nuclear and cytoplasmic in quiescent cells. AKT- or RSK-mediated phosphorylation on Thr-198, binds 14-3-3, translocates to the cytoplasm and promotes cell cycle progression. Mitogen-activated UHMK1 phosphorylation on Ser-10 also results in translocation to the cytoplasm and cell cycle progression. Phosphorylation on Ser-10 facilitates nuclear export. Translocates to the nucleus on phosphorylation of Tyr-88 and Tyr-89. Colocalizes at the endosome with SNX6; this leads to lysosomal degradation (By similarity)

**Tissue Location**

Expressed in kidney (at protein level) (PubMed:15509543). Expressed in all tissues tested (PubMed:8033212) Highest levels in skeletal muscle, lowest in liver and kidney (PubMed:8033212).


**Volume**

250 µl

**CDKN1B / p27 Kip1 Antibody (C-Terminus) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**CDKN1B / p27 Kip1 Antibody (C-Terminus) - Images**

Anti-CDKN1B / p27 Kip1 antibody IHC of human breast.

**CDKN1B / p27 Kip1 Antibody (C-Terminus) - Background**

Important regulator of cell cycle progression. Involved in G1 arrest. Potent inhibitor of cyclin E- and cyclin A-CDK2 complexes. Forms a complex with cyclin type D-CDK4 complexes and is involved in the assembly, stability, and modulation of CCND1- CDK4 complex activation. Acts either as an inhibitor or an activator of cyclin type D-CDK4 complexes depending on its phosphorylation state and/or stoichiometry.

**CDKN1B / p27 Kip1 Antibody (C-Terminus) - References**

Polyak K., et al. Cell 78:59-66(1994).  
Pietenpol J.A., et al. Cancer Res. 55:1206-1210(1995).  
Kalnina N., et al. Submitted (OCT-2004) to the EMBL/GenBank/DDBJ databases.  
Montagnoli A., et al. Genes Dev. 13:1181-1189(1999).  
Ishida N., et al. J. Biol. Chem. 275:25146-25154(2000).