

**FBXO11 Antibody (clone 4C12)**  
**Mouse Monoclonal Antibody**  
**Catalog # ALS14105****Specification**

---

**FBXO11 Antibody (clone 4C12) - Product Information**

Application	IF, IHC
Primary Accession	<a href="#">Q86XK2</a>
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Calculated MW	104kDa KDa

**FBXO11 Antibody (clone 4C12) - Additional Information****Gene ID** 80204**Other Names**

F-box only protein 11, Protein arginine N-methyltransferase 9, Vitiligo-associated protein 1, VIT-1, FBXO11, FBX11, PRMT9, VIT1

**Target/Specificity**

Human FBXO11

**Reconstitution & Storage**

Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles.

**Precautions**

FBXO11 Antibody (clone 4C12) is for research use only and not for use in diagnostic or therapeutic procedures.

**FBXO11 Antibody (clone 4C12) - Protein Information****Name** FBXO11**Synonyms** FBX11, PRMT9, VIT1**Function**

Substrate recognition component of a SCF (SKP1-CUL1-F-box protein) E3 ubiquitin-protein ligase complex which mediates the ubiquitination and subsequent proteasomal degradation of target proteins, such as DTL/CDT2, BCL6 and PRDM1/BLIMP1. The SCF(FBXO11) complex mediates ubiquitination and degradation of BCL6, thereby playing a role in the germinal center B-cells terminal differentiation toward memory B-cells and plasma cells. The SCF(FBXO11) complex also mediates ubiquitination and degradation of DTL, an important step for the regulation of TGF-beta signaling, cell migration and the timing of the cell-cycle progression and exit. Binds to and neddylates phosphorylated p53/TP53, inhibiting its transcriptional activity. Plays a role in the regulation of erythropoiesis but not myelopoiesis or megakaryopoiesis. Mechanistically, activates erythroid genes by mediating the degradation of BAHD1, a heterochromatin-associated protein

that recruits corepressors to H3K27me3 marks (PubMed:<a href="http://www.uniprot.org/citations/33156908" target="\_blank">33156908</a>). Participates in macrophage cell death and inflammation in response to bacterial toxins by regulating the expression of complement 5a receptor 1/C5AR1 and IL-1beta (PubMed:<a href="http://www.uniprot.org/citations/33156908" target="\_blank">33156908</a>). Acts as a critical regulator to determine the level of MHC-II by mediating the recognition of degran at the P/S/T domain of CIITA leading to its ubiquitination and subsequent degradation via the proteasome (PubMed:<a href="http://www.uniprot.org/citations/37279268" target="\_blank">37279268</a>). Participates in the antiviral response by initiating the activation of TBK1-IRF3-IFN-I axis. Mediates the 'Lys-63'-linked ubiquitination of TRAF3 to strengthen the interaction between TRAF3 and TBK1 (PubMed:<a href="http://www.uniprot.org/citations/36897010" target="\_blank">36897010</a>).

#### Cellular Location

Nucleus. Chromosome.

#### Tissue Location


Isoform 5 is expressed in keratinocytes, fibroblasts and melanocytes.

### FBXO11 Antibody (clone 4C12) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### FBXO11 Antibody (clone 4C12) - Images



Immunofluorescence of monoclonal antibody to FBXO11 on HeLa cell. [antibody concentration 10 ug/ml].



Anti-FBXO11 antibody IHC of human skin.

#### **FBXO11 Antibody (clone 4C12) - Background**

Substrate recognition component of a SCF (SKP1-CUL1-F-box protein) E3 ubiquitin-protein ligase complex which mediates the ubiquitination and subsequent proteasomal degradation of target proteins, such as DTL/CDT2, BCL6 and PRDM1/BLIMP1. The SCF(FBXO11) complex mediates ubiquitination and degradation of BCL6, thereby playing a role in the germinal center B-cells terminal differentiation toward memory B-cells and plasma cells. The SCF(FBXO11) complex also mediates ubiquitination and degradation of DTL, an important step for the regulation of TGF- $\beta$  signaling, cell migration and the timing of the cell-cycle progression and exit. Binds to and neddylates phosphorylated p53/TP53, inhibiting its transcriptional activity. SCF(FBXO11) does not seem to direct ubiquitination of p53/TP53.

#### **FBXO11 Antibody (clone 4C12) - References**

Abida W.M., et al. J. Biol. Chem. 282:1797-1804(2007).  
Mao Y.-M., et al. Submitted (FEB-2001) to the EMBL/GenBank/DDBJ databases.  
Totoki Y., et al. Submitted (APR-2005) to the EMBL/GenBank/DDBJ databases.  
Ota T., et al. Nat. Genet. 36:40-45(2004).  
Hillier L.W., et al. Nature 434:724-731(2005).