

PROKR2/Prokineticin Receptor 2 Antibody (Internal)
Goat Polyclonal Antibody
Catalog # ALS14256**Specification**

PROKR2/Prokineticin Receptor 2 Antibody (Internal) - Product Information

Application	WB, IHC
Primary Accession	Q8NFJ6
Reactivity	Human, Rat
Host	Goat
Clonality	Polyclonal
Calculated MW	44kDa KDa

PROKR2/Prokineticin Receptor 2 Antibody (Internal) - Additional Information**Gene ID** 128674**Other Names**

Prokineticin receptor 2, PK-R2, G-protein coupled receptor 73-like 1, GPR73b, GPRg2, PROKR2, GPR73L1, PKR2

Target/Specificity

Human PROKR2.

Reconstitution & Storage

Store at -20°C. Minimize freezing and thawing.

Precautions

PROKR2/Prokineticin Receptor 2 Antibody (Internal) is for research use only and not for use in diagnostic or therapeutic procedures.

PROKR2/Prokineticin Receptor 2 Antibody (Internal) - Protein Information**Name** PROKR2**Synonyms** GPR73L1, PKR2**Function**

Receptor for prokineticin 2. Exclusively coupled to the G(q) subclass of heteromeric G proteins. Activation leads to mobilization of calcium, stimulation of phosphoinositide turnover and activation of p44/p42 mitogen-activated protein kinase.

Cellular Location

Cell membrane; Multi-pass membrane protein

Tissue Location


Expressed in the ileocecum, thyroid gland, pituitary gland, salivary gland, adrenal gland, testis, ovary and brain

PROKR2/Prokineticin Receptor 2 Antibody (Internal) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PROKR2/Prokineticin Receptor 2 Antibody (Internal) - Images



PROKR2 antibody (1 ug/ml) staining of Rat Stomach lysate (35 ug protein/ml in RIPA buffer).



Anti-PROKR2 antibody IHC of human brain, cortex.

PROKR2/Prokineticin Receptor 2 Antibody (Internal) - Background

Receptor for prokineticin 2. Exclusively coupled to the G(q) subclass of heteromeric G proteins. Activation leads to mobilization of calcium, stimulation of phosphoinositide turnover and activation of p44/p42 mitogen-activated protein kinase.

PROKR2/Prokineticin Receptor 2 Antibody (Internal) - References

Lin D.C.-H., et al. J. Biol. Chem. 277:19276-19280(2002).

Soga T.,et al.Biochim. Biophys. Acta 1579:173-179(2002).
Martin A.L.,et al.Submitted (APR-2007) to the EMBL/GenBank/DDBJ databases.
Deloukas P.,et al.Nature 414:865-871(2001).
Marsango S.,et al.Cell. Mol. Life Sci. 68:2919-2929(2011).