

PRDX6 / Peroxiredoxin 6 Antibody
Rabbit Polyclonal Antibody
Catalog # ALS14260**Specification**

PRDX6 / Peroxiredoxin 6 Antibody - Product Information

Application	WB, IHC
Primary Accession	P30041
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	25kDa KDa

PRDX6 / Peroxiredoxin 6 Antibody - Additional Information**Gene ID** 9588**Other Names**

Peroxiredoxin-6, 1.11.1.15, 1-Cys peroxiredoxin, 1-Cys PRX, 24 kDa protein, Acidic calcium-independent phospholipase A2, aiPLA2, 3.1.1.-, Antioxidant protein 2, Liver 2D page spot 40, Non-selenium glutathione peroxidase, NSGPx, 1.11.1.9, Red blood cells page spot 12, PRDX6, AOP2, KIAA0106

Reconstitution & Storage

May be stored at 4°C for short-term only. For long-term storage and to avoid repeated freezing and thawing, aliquot. Store at -20°C. Aliquots are stable for at least 12 months at -20°C

Precautions

PRDX6 / Peroxiredoxin 6 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

PRDX6 / Peroxiredoxin 6 Antibody - Protein Information**Name** PRDX6**Synonyms** AOP2, KIAA0106**Function**

Thiol-specific peroxidase that catalyzes the reduction of hydrogen peroxide and organic hydroperoxides to water and alcohols, respectively (PubMed:9497358, PubMed:10893423). Can reduce H₂O₂ and short chain organic, fatty acid, and phospholipid hydroperoxides (PubMed:10893423). Also has phospholipase activity, can therefore either reduce the oxidized sn-2 fatty acyl group of phospholipids (peroxidase activity) or hydrolyze the sn-2 ester bond of phospholipids (phospholipase activity) (PubMed:10893423, PubMed:26830860).

target="_blank">26830860). These activities are dependent on binding to phospholipids at acidic pH and to oxidized phospholipids at cytosolic pH (PubMed:10893423). Plays a role in cell protection against oxidative stress by detoxifying peroxides and in phospholipid homeostasis (PubMed:10893423). Exhibits acyl-CoA-dependent lysophospholipid acyltransferase which mediates the conversion of lysophosphatidylcholine (1-acyl-sn-glycero-3-phosphocholine or LPC) into phosphatidylcholine (1,2-diacyl-sn-glycero-3-phosphocholine or PC) (PubMed:26830860). Shows a clear preference for LPC as the lysophospholipid and for palmitoyl CoA as the fatty acyl substrate (PubMed:26830860).

Cellular Location

Cytoplasm. Lysosome {ECO:0000250|UniProtKB:O35244}. Note=Also found in lung secretory organelles (lamellar bodies). {ECO:0000250|UniProtKB:O35244}

Volume


50 µl


PRDX6 / Peroxiredoxin 6 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PRDX6 / Peroxiredoxin 6 Antibody - Images





Anti-PRDX6 / Peroxiredoxin 6 antibody IHC of human adrenal.



Anti-PRDX6 / Peroxiredoxin 6 antibody IHC of human testis.

PRDX6 / Peroxiredoxin 6 Antibody - Background

Involved in redox regulation of the cell. Can reduce H_2O_2 and short chain organic, fatty acid, and phospholipid hydroperoxides. May play a role in the regulation of phospholipid turnover as well as in protection against oxidative injury.

PRDX6 / Peroxiredoxin 6 Antibody - References

- Kim T.-S.,et al.J. Biol. Chem. 272:2542-2550(1997).
- Kim T.-S.,et al.J. Biol. Chem. 272:10981-10981(1997).
- Frank S.,et al.Oncogene 14:915-921(1997).
- Nagase T.,et al.DNA Res. 2:37-43(1995).
- Ota T.,et al.Nat. Genet. 36:40-45(2004).