

KCNC2 / Kv3.2 Antibody (aa589-638)
Rabbit Polyclonal Antibody
Catalog # ALS14296**Specification****KCNC2 / Kv3.2 Antibody (aa589-638) - Product Information**

Application	WB, IHC
Primary Accession	Q96PR1
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	70kDa KDa

KCNC2 / Kv3.2 Antibody (aa589-638) - Additional Information**Gene ID** 3747**Other Names**

Potassium voltage-gated channel subfamily C member 2, Voltage-gated potassium channel Kv3.2, KCNC2

Target/Specificity

Potassium Channel Kv3.2b Antibody detects endogenous levels of total Potassium Channel Kv3.2b protein.

Reconstitution & Storage

Store at -20°C.

Precautions

KCNC2 / Kv3.2 Antibody (aa589-638) is for research use only and not for use in diagnostic or therapeutic procedures.

KCNC2 / Kv3.2 Antibody (aa589-638) - Protein Information**Name** KCNC2 ([HGNC:6234](#))**Function**

Voltage-gated potassium channel that mediates transmembrane potassium transport in excitable membranes, primarily in the brain. Contributes to the regulation of the fast action potential repolarization and in sustained high-frequency firing in neurons of the central nervous system. Homotetramer channels mediate delayed-rectifier voltage-dependent potassium currents that activate rapidly at high- threshold voltages and inactivate slowly. Forms tetrameric channels through which potassium ions pass in accordance with their electrochemical gradient. The channel alternates between opened and closed conformations in response to the voltage difference across the membrane (Probable) (PubMed: [15709110](http://www.uniprot.org/citations/15709110) target="_blank">15709110, PubMed: [35314505](http://www.uniprot.org/citations/35314505) target="_blank">35314505, PubMed: [36090251](http://www.uniprot.org/citations/36090251) target="_blank">36090251). Can form functional homotetrameric and heterotetrameric

channels that contain variable proportions of KCNC1, and possibly other family members as well; channel properties depend on the type of alpha subunits that are part of the channel. Channel properties may be modulated either by the association with ancillary subunits, such as KCNE1, KCNE2 or KCNE3 or indirectly by nitric oxide (NO) through a cGMP- and PKG-mediated signaling cascade, slowing channel activation and deactivation of delayed rectifier potassium channels (By similarity). Contributes to fire sustained trains of very brief action potentials at high frequency in retinal ganglion cells, thalamocortical and suprachiasmatic nucleus (SCN) neurons and in hippocampal and neocortical interneurons (PubMed:15709110). Sustained maximal action potential firing frequency in inhibitory hippocampal interneurons is negatively modulated by histamine H2 receptor activation in a cAMP- and protein kinase (PKA) phosphorylation- dependent manner. Plays a role in maintaining the fidelity of synaptic transmission in neocortical GABAergic interneurons by generating action potential (AP) repolarization at nerve terminals, thus reducing spike- evoked calcium influx and GABA neurotransmitter release. Required for long-range synchronization of gamma oscillations over distance in the neocortex. Contributes to the modulation of the circadian rhythm of spontaneous action potential firing in suprachiasmatic nucleus (SCN) neurons in a light-dependent manner (By similarity).

Cellular Location

Cell membrane {ECO:0000250|UniProtKB:Q14B80}; Multi-pass membrane protein. Membrane {ECO:0000250|UniProtKB:Q14B80}; Multi-pass membrane protein. Perikaryon {ECO:0000250|UniProtKB:Q14B80}. Cell projection, axon {ECO:0000250|UniProtKB:Q14B80}. Cell projection, dendrite {ECO:0000250|UniProtKB:Q14B80}. Postsynaptic cell membrane {ECO:0000250|UniProtKB:Q14B80}. Presynaptic cell membrane {ECO:0000250|UniProtKB:Q14B80}. Synapse, synaptosome {ECO:0000250|UniProtKB:P22462}. Synapse {ECO:0000250|UniProtKB:P22462}. Apical cell membrane {ECO:0000250|UniProtKB:P22462}. Basolateral cell membrane {ECO:0000250|UniProtKB:P22462}. Note=Colocalizes with parvalbumin in globus pallidus neurons. Localizes in thalamocortical axons and synapses. Localizes on the surface of cell somata, proximal dendrites and axonal membranes. Also detected throughout the neuropil Localized in starburst cell somata and proximal dendrite processes Colocalized with GABA in presynaptic terminals. Clustered in patches in somatic and proximal dendritic membrane as well as in axons and presynaptic terminals of GABAergic interneurons; some of these patches are found near postsynaptic sites. {ECO:0000250|UniProtKB:P22462, ECO:0000250|UniProtKB:Q14B80}

Volume


50 µl


KCNC2 / Kv3.2 Antibody (aa589-638) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

KCNC2 / Kv3.2 Antibody (aa589-638) - Images





Western blot of extracts from HepG2 cells, using KCNC2 antibody.



Anti-KCNC2 / Kv3.2 antibody IHC of human brain, cortex.

KCNC2 / Kv3.2 Antibody (aa589-638) - Background

Mediates the voltage-dependent potassium ion permeability of excitable membranes. Assuming opened or closed conformations in response to the voltage difference across the membrane, the protein forms a potassium-selective channel through which potassium ions may pass in accordance with their electrochemical gradient. Channel properties are modulated by subunit assembly (By similarity).

KCNC2 / Kv3.2 Antibody (aa589-638) - References

- Haas M.,et al.Mamm. Genome 4:711-715(1993).
- Isbrandt D.,et al.Submitted (JUN-2002) to the EMBL/GenBank/DDBJ databases.
- Ota T.,et al.Nat. Genet. 36:40-45(2004).
- Nakajima D.,et al.Submitted (MAR-2005) to the EMBL/GenBank/DDBJ databases.
- Scherer S.E.,et al.Nature 440:346-351(2006).