

SLC9A3R1 / NHERF1 / EBP50 Antibody (clone 2B7)

Mouse Monoclonal Antibody Catalog # ALS14426

Specification

SLC9A3R1 / NHERF1 / EBP50 Antibody (clone 2B7) - Product Information

Application IHC
Primary Accession O14745
Reactivity Human
Host Mouse
Clonality Monoclonal
Calculated MW 39kDa KDa

SLC9A3R1 / NHERF1 / EBP50 Antibody (clone 2B7) - Additional Information

Gene ID 9368

Other Names

Na(+)/H(+) exchange regulatory cofactor NHE-RF1, NHERF-1, Ezrin-radixin-moesin-binding phosphoprotein 50, EBP50, Regulatory cofactor of Na(+)/H(+) exchanger, Sodium-hydrogen exchanger regulatory factor 1, Solute carrier family 9 isoform A3 regulatory factor 1, SLC9A3R1, NHERF, NHERF1

Target/Specificity

Human EBP50

Reconstitution & Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

SLC9A3R1 / NHERF1 / EBP50 Antibody (clone 2B7) is for research use only and not for use in diagnostic or therapeutic procedures.

SLC9A3R1 / NHERF1 / EBP50 Antibody (clone 2B7) - Protein Information

Name NHERF1 (HGNC:11075)

Synonyms NHERF, SLC9A3R1

Function

Scaffold protein that connects plasma membrane proteins with members of the ezrin/moesin/radixin family and thereby helps to link them to the actin cytoskeleton and to regulate their surface expression. Necessary for recycling of internalized ADRB2. Was first known to play a role in the regulation of the activity and subcellular location of SLC9A3. Necessary for cAMP-mediated phosphorylation and inhibition of SLC9A3. May enhance Wnt signaling. May participate in HTR4 targeting to microvilli (By similarity). Involved in the regulation of phosphate reabsorption in the renal proximal tubules. Involved in sperm capacitation. May participate in the regulation of the chloride and bicarbonate homeostasis in spermatozoa.



Cellular Location

Cytoplasm. Apical cell membrane. Endomembrane system; Peripheral membrane protein. Cell projection, filopodium. Cell projection, ruffle. Cell projection, microvillus. Note=Translocates from the cytoplasm to the apical cell membrane in a PODXL-dependent manner. Colocalizes with CFTR at the midpiece of sperm tail (By similarity). Colocalizes with actin in microvilli-rich apical regions of the syncytiotrophoblast. Found in microvilli, ruffling membrane and filopodia of HeLa cells. Present in lipid rafts of T-cells.

Tissue Location


Detected in liver, kidney, pancreas, prostate, spleen, small intestine and placenta, in particular in the syncytiotrophoblast.

SLC9A3R1 / NHERF1 / EBP50 Antibody (clone 2B7) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

SLC9A3R1 / NHERF1 / EBP50 Antibody (clone 2B7) - Images




Anti-SLC9A3R1 / NHERF1 antibody IHC of human small intestine.

SLC9A3R1 / NHERF1 / EBP50 Antibody (clone 2B7) - Background

Scaffold protein that connects plasma membrane proteins with members of the ezrin/moesin/radixin family and thereby helps to link them to the actin cytoskeleton and to regulate their surface expression. Necessary for recycling of internalized ADRB2. Was first known to play a role in the regulation of the activity and subcellular location of SLC9A3. Necessary for cAMP-mediated phosphorylation and inhibition of SLC9A3. May enhance Wnt signaling. May participate in HTR4 targeting to microvilli (By similarity). Involved in the regulation of phosphate reabsorption in the renal proximal tubules. Involved in sperm capacitation. May participate in the regulation of the chloride and bicarbonate homeostasis in spermatozoa.

SLC9A3R1 / NHERF1 / EBP50 Antibody (clone 2B7) - References





Reczek D.,et al.J. Cell Biol. 139:169-179(1997).

Murthy A.,et al.J. Biol. Chem. 273:1273-1276(1998).

Ota T.,et al.Nat. Genet. 36:40-45(2004).

Zody M.C.,et al.Nature 440:1045-1049(2006).

Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.