

TLR1 Antibody (aa400-450)
Rabbit Polyclonal Antibody
Catalog # ALS14593**Specification**

TLR1 Antibody (aa400-450) - Product Information

Application	WB
Primary Accession	Q15399
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	90kDa KDa

TLR1 Antibody (aa400-450) - Additional Information**Gene ID** 7096**Other Names**

Toll-like receptor 1, Toll/interleukin-1 receptor-like protein, TIL, CD281, TLR1, KIAA0012

Target/Specificity

A KLH-conjugated synthetic peptide corresponding to a sequence within amino acids 400-450 of human TLR1.

Reconstitution & Storage

Long term: -20°C; Short term: +4°C; Avoid freeze-thaw cycles.

Precautions

TLR1 Antibody (aa400-450) is for research use only and not for use in diagnostic or therapeutic procedures.

TLR1 Antibody (aa400-450) - Protein Information**Name** TLR1**Synonyms** KIAA0012**Function**

Participates in the innate immune response to microbial agents. Specifically recognizes diacylated and triacylated lipopeptides. Cooperates with TLR2 to mediate the innate immune response to bacterial lipoproteins or lipopeptides (PubMed:21078852). Forms the activation cluster TLR2:TLR1:CD14 in response to triacylated lipopeptides, this cluster triggers signaling from the cell surface and subsequently is targeted to the Golgi in a lipid-raft dependent pathway (PubMed:16880211). Acts via MYD88 and TRAF6, leading to NF-kappa-B activation, cytokine secretion and the inflammatory response.

Cellular Location

Cell membrane; Single-pass type I membrane protein. Cytoplasmic vesicle, phagosome membrane {ECO:0000250|UniProtKB:Q9EPQ1}; Single-pass type I membrane protein. Membrane raft. Golgi apparatus. Note=Does not reside in lipid rafts before stimulation but accumulates increasingly in the raft upon the presence of the microbial ligand. In response to triacylated lipoproteins, TLR2:TLR1 heterodimers are recruited in lipid rafts, this recruitment determine the intracellular targeting to the Golgi apparatus.

Tissue Location


Ubiquitous. Highly expressed in spleen, ovary, peripheral blood leukocytes, thymus and small intestine

TLR1 Antibody (aa400-450) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TLR1 Antibody (aa400-450) - Images



Western blot of TLR1 in Ramos (A), Raw (B), and (C) HEK293 cells transfected with full length...

TLR1 Antibody (aa400-450) - Background

Participates in the innate immune response to microbial agents. Specifically recognizes diacylated and triacylated lipopeptides. Cooperates with TLR2 to mediate the innate immune response to bacterial lipoproteins or lipopeptides. Acts via MYD88 and TRAF6, leading to NF-kappa-B activation, cytokine secretion and the inflammatory response (By similarity).

TLR1 Antibody (aa400-450) - References

Rock F.L.,et al.Proc. Natl. Acad. Sci. U.S.A. 95:588-593(1998).
Nakajima T.,et al.Immunogenetics 60:727-735(2008).
Georgel P.,et al.PLoS ONE 4:E7803-E7803(2009).
Nomura N.,et al.DNA Res. 1:27-35(1994).
Wiemann S.,et al.Genome Res. 11:422-435(2001).