

NOD1 Antibody (aa930-949)
Rabbit Polyclonal Antibody
Catalog # ALS14598**Specification**

NOD1 Antibody (aa930-949) - Product Information

Application	WB, IHC
Primary Accession	O9Y239
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	108kDa KDa

NOD1 Antibody (aa930-949) - Additional Information**Gene ID** 10392**Other Names**

Nucleotide-binding oligomerization domain-containing protein 1, Caspase recruitment domain-containing protein 4, NOD1, CARD4

Target/Specificity

Amino acids 930-949 of human Nod1 protein were used as the immunogen.

Reconstitution & Storage

Long term: -20°C; Short term: +4°C; Avoid freeze-thaw cycles.

Precautions

NOD1 Antibody (aa930-949) is for research use only and not for use in diagnostic or therapeutic procedures.

NOD1 Antibody (aa930-949) - Protein Information**Name** NOD1 {ECO:0000303|PubMed:10329646, ECO:0000312|HGNC:HGNC:16390}**Function**

Pattern recognition receptor (PRR) that detects bacterial peptidoglycan fragments and other danger signals and thus participates in both innate and adaptive immune responses (PubMed:11058605, PubMed:12796777, PubMed:12791997, PubMed:15044951, PubMed:16172124, PubMed:19043560, PubMed:22672233, PubMed:27099311). Specifically recognizes and binds gamma-D-glutamyl-meso-diaminopimelic acid (iE- DAP), a dipeptide present in peptidoglycan of Gram-negative bacteria (PubMed:<a

<http://www.uniprot.org/citations/12871942> target="_blank">12871942, PubMed:12796777, PubMed:12791997, PubMed:16211083, PubMed:16172124). Preferentially binds iE-DAP in tripeptide-containing mucopeptides (MurNAc-TriDAP or TriDAP) (PubMed:16211083). Ligand binding triggers oligomerization that facilitates the binding and subsequent activation of the proximal adapter receptor-interacting RIPK2 (PubMed:12796777, PubMed:12791997, PubMed:17054981). Following recruitment, RIPK2 undergoes 'Met-1'- (linear) and 'Lys-63'-linked polyubiquitination by E3 ubiquitin-protein ligases XIAP, BIRC2, BIRC3 and the LUBAC complex, becoming a scaffolding protein for downstream effectors, triggering activation of the NF-kappa-B and MAP kinases signaling (PubMed:10880512, PubMed:12791997, PubMed:19043560). This in turn leads to the transcriptional activation of hundreds of genes involved in immune response (PubMed:10880512, PubMed:19043560). Also acts as a regulator of antiviral response elicited by dsRNA and the expression of RLR pathway members by targeting IFIH1 and TRAF3 to modulate the formation of IFIH1-MAVS and TRAF3-MAVS complexes leading to increased transcription of type I IFNs (PubMed:32169843). Also acts as a regulator of autophagy via its interaction with ATG16L1, possibly by recruiting ATG16L1 at the site of bacterial entry (By similarity). Besides recognizing pathogens, also involved in the endoplasmic reticulum stress response: acts by sensing and binding to the cytosolic metabolite sphingosine-1-phosphate generated in response to endoplasmic reticulum stress, initiating an inflammation process that leads to activation of the NF-kappa-B and MAP kinases signaling (PubMed:27007849, PubMed:33942347). In addition, plays a role in insulin trafficking in beta cells in a cell-autonomous manner (By similarity). Mechanistically, upon recognizing cognate ligands, NOD1 and RIPK2 localize to insulin vesicles where they recruit RAB1A to direct insulin trafficking through the cytoplasm (By similarity).

Cellular Location

Cell membrane; Lipid-anchor. Apical cell membrane. Basolateral cell membrane. Cytoplasm. Note=Detected in the cytoplasm and at the cell membrane (PubMed:31649195). Following bacterial infection, localizes to bacterial entry sites in the cell membrane (PubMed:31649195). Recruited to the basolateral and apical membranes in polarized epithelial cells (PubMed:19043560)

Tissue Location

Highly expressed in adult heart, skeletal muscle, pancreas, spleen and ovary (PubMed:10224040). Also detected in placenta, lung, liver, kidney, thymus, testis, small intestine and colon (PubMed:10224040).

Volume


50 µl

NOD1 Antibody (aa930-949) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NOD1 Antibody (aa930-949) - Images



Western blot of NOD1 in (A) recombinant fusion protein containing amino acids 930-949 and (B)...



Anti-NOD1 antibody IHC of human pancreas.

NOD1 Antibody (aa930-949) - Background

Enhances caspase-9-mediated apoptosis. Induces NF-kappa-B activity via RIPK2 and IKK-gamma. Confers responsiveness to intracellular bacterial lipopolysaccharides (LPS). Forms an intracellular sensing system along with ARHGEF2 for the detection of microbial effectors during cell invasion by pathogens. Required for RHOA and RIPK2 dependent NF-kappa-B signaling pathway activation upon *S.flexneri* cell invasion. Involved not only in sensing peptidoglycan (PGN)-derived muropeptides but also in the activation of NF-kappa-B by *Shigella* effector proteins IpgB2 and OspB. Recruits NLRP10 to the cell membrane following bacterial infection.

NOD1 Antibody (aa930-949) - References

Bertin J., et al. J. Biol. Chem. 274:12955-12958(1999).

Inohara N.,et al.J. Biol. Chem. 274:14560-14567(1999).

Ota T.,et al.Nat. Genet. 36:40-45(2004).

Hillier L.W.,et al.Nature 424:157-164(2003).

Inohara N.,et al.J. Biol. Chem. 276:2551-2554(2001).