

**AURKA / Aurora-A Antibody (aa100-150)**  
**Rabbit Polyclonal Antibody**  
**Catalog # ALS14602****Specification**

---

**AURKA / Aurora-A Antibody (aa100-150) - Product Information**

|                   |                        |
|-------------------|------------------------|
| Application       | WB                     |
| Primary Accession | <a href="#">O14965</a> |
| Reactivity        | Human, Mouse, Rat      |
| Host              | Rabbit                 |
| Clonality         | Polyclonal             |
| Calculated MW     | 46kDa KDa              |

**AURKA / Aurora-A Antibody (aa100-150) - Additional Information****Gene ID** 6790**Other Names**

Aurora kinase A, 2.7.11.1, Aurora 2, Aurora/IPL1-related kinase 1, ARK-1, Aurora-related kinase 1, hARK1, Breast tumor-amplified kinase, Serine/threonine-protein kinase 15, Serine/threonine-protein kinase 6, Serine/threonine-protein kinase aurora-A, AURKA

**Target/Specificity**

A portion of amino acids 100-150 of human AIK1 was used as the immunogen.

**Reconstitution & Storage**

Long term: -20°C; Short term: +4°C; Avoid freeze-thaw cycles.

**Precautions**

AURKA / Aurora-A Antibody (aa100-150) is for research use only and not for use in diagnostic or therapeutic procedures.

**AURKA / Aurora-A Antibody (aa100-150) - Protein Information****Name** AURKA ([HGNC:11393](#))**Function**

Mitotic serine/threonine kinase that contributes to the regulation of cell cycle progression (PubMed:<a href="http://www.uniprot.org/citations/26246606" target="\_blank">26246606</a>, PubMed:<a href="http://www.uniprot.org/citations/12390251" target="\_blank">12390251</a>, PubMed:<a href="http://www.uniprot.org/citations/18615013" target="\_blank">18615013</a>, PubMed:<a href="http://www.uniprot.org/citations/11039908" target="\_blank">11039908</a>, PubMed:<a href="http://www.uniprot.org/citations/17125279" target="\_blank">17125279</a>, PubMed:<a href="http://www.uniprot.org/citations/17360485" target="\_blank">17360485</a>). Associates with the centrosome and the spindle microtubules during mitosis and plays a critical role in various mitotic events including the establishment of mitotic spindle, centrosome duplication, centrosome separation as well as maturation, chromosomal alignment, spindle assembly checkpoint, and cytokinesis (PubMed:<a

[26246606](http://www.uniprot.org/citations/26246606), PubMed: [14523000](http://www.uniprot.org/citations/14523000)). Required for normal spindle positioning during mitosis and for the localization of NUMA1 and DCTN1 to the cell cortex during metaphase (PubMed: [27335426](http://www.uniprot.org/citations/27335426)). Required for initial activation of CDK1 at centrosomes (PubMed: [13678582](http://www.uniprot.org/citations/13678582), PubMed: [15128871](http://www.uniprot.org/citations/15128871)). Phosphorylates numerous target proteins, including ARHGEF2, BORA, BRCA1, CDC25B, DLGP5, HDAC6, KIF2A, LATS2, NDEL1, PARD3, PPP1R2, PLK1, RASSF1, TACC3, p53/TP53 and TPX2 (PubMed: [18056443](http://www.uniprot.org/citations/18056443), PubMed: [15128871](http://www.uniprot.org/citations/15128871), PubMed: [14702041](http://www.uniprot.org/citations/14702041), PubMed: [11551964](http://www.uniprot.org/citations/11551964), PubMed: [15147269](http://www.uniprot.org/citations/15147269), PubMed: [15987997](http://www.uniprot.org/citations/15987997), PubMed: [17604723](http://www.uniprot.org/citations/17604723), PubMed: [18615013](http://www.uniprot.org/citations/18615013)). Regulates KIF2A tubulin depolymerase activity (PubMed: [19351716](http://www.uniprot.org/citations/19351716)). Important for microtubule formation and/or stabilization (PubMed: [18056443](http://www.uniprot.org/citations/18056443)). Required for normal axon formation (PubMed: [19812038](http://www.uniprot.org/citations/19812038)). Plays a role in microtubule remodeling during neurite extension (PubMed: [19668197](http://www.uniprot.org/citations/19668197)). Also acts as a key regulatory component of the p53/TP53 pathway, and particularly the checkpoint- response pathways critical for oncogenic transformation of cells, by phosphorylating and destabilizing p53/TP53 (PubMed: [14702041](http://www.uniprot.org/citations/14702041)). Phosphorylates its own inhibitors, the protein phosphatase type 1 (PP1) isoforms, to inhibit their activity (PubMed: [11551964](http://www.uniprot.org/citations/11551964)). Inhibits cilia outgrowth (By similarity). Required for cilia disassembly via phosphorylation of HDAC6 and subsequent deacetylation of alpha-tubulin (PubMed: [17604723](http://www.uniprot.org/citations/17604723), PubMed: [20643351](http://www.uniprot.org/citations/20643351)). Regulates protein levels of the anti-apoptosis protein BIRC5 by suppressing the expression of the SCF(FBXL7) E3 ubiquitin-protein ligase substrate adapter FBXL7 through the phosphorylation of the transcription factor FOXP1 (PubMed: [28218735](http://www.uniprot.org/citations/28218735)).

### Cellular Location

Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, spindle pole. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome, centriole {ECO:0000250|UniProtKB:P97477}. Cell projection, neuron projection {ECO:0000250|UniProtKB:P97477}. Cell projection, cilium. Cytoplasm, cytoskeleton, cilium basal body. Basolateral cell membrane {ECO:0000250|UniProtKB:F1PNY0}. Note=Detected at the neurite hillock in developing neurons (By similarity). Localizes at the centrosome in mitotic cells from early prophase until telophase, but also localizes to the spindle pole MTs from prophase to anaphase (PubMed:9606188, PubMed:17229885, PubMed:21225229). Colocalized with SIRT2 at centrosome (PubMed:22014574). Moves to the midbody during both telophase and cytokinesis (PubMed:17726514). Associates with both the pericentriolar material (PCM) and centrioles (PubMed:22014574). The localization to the spindle poles is regulated by AAAS (PubMed:26246606) {ECO:0000250|UniProtKB:P97477, ECO:0000269|PubMed:17229885, ECO:0000269|PubMed:17726514, ECO:0000269|PubMed:21225229, ECO:0000269|PubMed:22014574, ECO:0000269|PubMed:26246606, ECO:0000269|PubMed:9606188}

### Tissue Location


Highly expressed in testis and weakly in skeletal muscle, thymus and spleen. Also highly expressed in colon, ovarian, prostate, neuroblastoma, breast and cervical cancer cell lines

### **AURKA / Aurora-A Antibody (aa100-150) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **AURKA / Aurora-A Antibody (aa100-150) - Images**



Western blot of AIK1 in A) HCT-116, B) human testis, C) mouse testis, and D) rat testis lysate...

### **AURKA / Aurora-A Antibody (aa100-150) - Background**

Mitotic serine/threonine kinases that contributes to the regulation of cell cycle progression. Associates with the centrosome and the spindle microtubules during mitosis and plays a critical role in various mitotic events including the establishment of mitotic spindle, centrosome duplication, centrosome separation as well as maturation, chromosomal alignment, spindle assembly checkpoint, and cytokinesis. Required for initial activation of CDK1 at centrosomes. Phosphorylates numerous target proteins, including ARHGEF2, BORA, BRCA1, CDC25B, DLGP5, HDAC6, KIF2A, LATS2, NDEL1, PARD3, PPP1R2, PLK1, RASSF1, TACC3, p53/TP53 and TPX2. Regulates KIF2A tubulin depolymerase activity. Required for normal axon formation. Plays a role in microtubule remodeling during neurite extension. Important for microtubule formation and/or stabilization. Also acts as a key regulatory component of the p53/TP53 pathway, and particularly the checkpoint-response pathways critical for oncogenic transformation of cells, by phosphorylating and stabilizing p53/TP53. Phosphorylates its own inhibitors, the protein phosphatase type 1 (PP1) isoforms, to inhibit their activity. Necessary for proper cilia disassembly prior to mitosis.

### **AURKA / Aurora-A Antibody (aa100-150) - References**

Kimura M.,et al.J. Biol. Chem. 272:13766-13771(1997).  
Shindo M.,et al.Biochem. Biophys. Res. Commun. 244:285-292(1998).  
Zhou H.,et al.Nat. Genet. 20:189-193(1998).  
Wang L.,et al.Submitted (OCT-1999) to the EMBL/GenBank/DDBJ databases.  
Deloukas P.,et al.Nature 414:865-871(2001).