

SUCNR1 / GPR91 Antibody (aa100-149)

Rabbit Polyclonal Antibody Catalog # ALS14625

Specification

SUCNR1 / GPR91 Antibody (aa100-149) - Product Information

Application WB, IHC Primary Accession Q9BXA5

Reactivity Human, Mouse, Rat

Host Rabbit
Clonality Polyclonal
Calculated MW 39kDa KDa

SUCNR1 / GPR91 Antibody (aa100-149) - Additional Information

Gene ID 56670

Other Names

Succinate receptor 1, G-protein coupled receptor 91, P2Y purinoceptor 1-like, SUCNR1, GPR91

Target/Specificity

SUCNR1 Antibody detects endogenous levels of total SUCNR1 protein.

Reconstitution & Storage

Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles.

Precautions

SUCNR1 / GPR91 Antibody (aa100-149) is for research use only and not for use in diagnostic or therapeutic procedures.

SUCNR1 / GPR91 Antibody (aa100-149) - Protein Information

Name SUCNR1 (HGNC:4542)

Synonyms GPR91

Function

G protein-coupled receptor for succinate able to mediate signaling through Gq/GNAQ or Gi/GNAI second messengers depending on the cell type and the processes regulated (By similarity) (PubMed:15141213, PubMed:34133934, PubMed:23770096). Succinate-SUCNR1 signaling serves as a link between metabolic stress, inflammation and energy homeostasis (PubMed:18820681, PubMed:34133934, PubMed:34133934, PubMed:34133934, PubMed:34133934, PubMed:<a href="http://www.uniprot.org/citations/18820681" targ



M2 macrophages versus M1 phenotype through Gq signaling by regulating the transcription of genes involoved in immune function (PubMed:34133934). In activated M1 macrophages, plays a pro-inflammatory role in response to LPS (By similarity). Expressed in dendritic cells, where it is involved in the sensing of immunological danger and enhances immunity. Mediates succinate triggered intracelleular calcium mobilization, induces migratory responses and acts in synergy with Toll-like receptor ligands for the production of proinflammatory cytokines as well as an enhancement of antigen-specific activation of helper T cells (PubMed:18820681). In the small intestine, mediates the activation of tuft cells by dietary succinate and triggers type 2 immunity (By similarity). In adipocytes, plays an important role in the control of energy metabolism. In response to succinate, controls leptin expression in an AMPK-JNK-CEBPA-dependent as well as circadian clock-regulated manner (By similarity). In muscle tissue, is expressed in non-muscle cells and coordinates muscle remodeling in response to the succinate produced during exercise training in a paracrine manner (By similarity). In retina, acts as a mediator of vessel growth during retinal development. In response to succinate, regulates the production of angiogenic factors, including VEGF, by retinal ganglion neurons (By similarity).

Cellular Location

Cell membrane; Multi-pass membrane protein

Tissue Location

Expressed specifically in kidney (PubMed:11273702). Highly expressed in immature dendritic cells, expression rapidly downregulates after maturation. Also expressed in macrophages (PubMed:18820681).

Volume


50 μl

SUCNR1 / GPR91 Antibody (aa100-149) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cvtometv
- Cell Culture


SUCNR1 / GPR91 Antibody (aa100-149) - Images







Western blot of extracts from HUVEC/MCF-7 cells, using SUCNR1 Antibody.



Anti-GPR91 antibody IHC of human adrenal cortex.

SUCNR1 / GPR91 Antibody (aa100-149) - Background

Receptor for succinate.

SUCNR1 / GPR91 Antibody (aa100-149) - References

Wittenberger T.,et al.J. Mol. Biol. 307:799-813(2001). Zhang W.,et al.Submitted (MAR-2000) to the EMBL/GenBank/DDBJ databases. Kaighin V.A.,et al.Submitted (DEC-2007) to the EMBL/GenBank/DDBJ databases. Ota T.,et al.Nat. Genet. 36:40-45(2004). Muzny D.M.,et al.Nature 440:1194-1198(2006).