

NDRG4 Antibody (clone 2G3)

Mouse Monoclonal Antibody Catalog # ALS14653

Specification

NDRG4 Antibody (clone 2G3) - Product Information

Application WB
Primary Accession O9ULPO

Reactivity Human, Mouse

Host Mouse
Clonality Monoclonal
Calculated MW 38kDa KDa

NDRG4 Antibody (clone 2G3) - Additional Information

Gene ID 65009

Other Names

Protein NDRG4, Brain development-related molecule 1, N-myc downstream-regulated gene 4 protein, Vascular smooth muscle cell-associated protein 8, SMAP-8, NDRG4, BDM1, KIAA1180

Target/Specificity

Human NDRG4

Reconstitution & Storage

Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles.

Precautions

NDRG4 Antibody (clone 2G3) is for research use only and not for use in diagnostic or therapeutic procedures.

NDRG4 Antibody (clone 2G3) - Protein Information

Name NDRG4

Synonyms BDM1, KIAA1180

Function

Contributes to the maintenance of intracerebral BDNF levels within the normal range, which is necessary for the preservation of spatial learning and the resistance to neuronal cell death caused by ischemic stress (By similarity). May enhance growth factor-induced ERK1 and ERK2 phosphorylation, including that induced by PDGF and FGF. May attenuate NGF-promoted ELK1 phosphorylation in a microtubule-dependent manner.

Cellular Location

Cytoplasm, cytosol

Tissue Location




Expressed predominantly in brain and heart (at protein level). In the brain, detected in astrocytes. Isoform 1 and isoform 2 are only expressed in brain. Isoform 3 is expressed in both heart and brain. Up-regulated in glioblastoma multiforme cells

NDRG4 Antibody (clone 2G3) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

NDRG4 Antibody (clone 2G3) - Images



Antibody ALS14653 Western blot of NDRG4 expression in Raw 264.7.

NDRG4 Antibody (clone 2G3) - Background

Contributes to the maintenance of intracerebral BDNF levels within the normal range, which is necessary for the preservation of spatial learning and the resistance to neuronal cell death caused by ischemic stress (By similarity). May enhance growth factor-induced ERK1 and ERK2 phosphorylation, including that induced by PDGF and FGF. May attenuate NGF-promoted ELK1 phosphorylation in a microtubule-dependent manner.

NDRG4 Antibody (clone 2G3) - References

Zhou R.-H.,et al.Genomics 73:86-97(2001). Qu X.,et al.Mol. Cell. Biochem. 229:35-44(2002). Nishimoto S.,et al.Eur. J. Biochem. 270:2521-2531(2003). Wiemann S.,et al.Genome Res. 11:422-435(2001). Ota T.,et al.Nat. Genet. 36:40-45(2004).