

**ATG9A Antibody (C-Terminus)**  
**Rabbit Polyclonal Antibody**  
**Catalog # ALS14712****Specification****ATG9A Antibody (C-Terminus) - Product Information**

Application	WB, IHC-P, IF
Primary Accession	<a href="#">Q7Z3C6</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	94kDa KDa
Dilution	WB~~1:1000 IHC-P~~N/A IF~~1:50~200

**ATG9A Antibody (C-Terminus) - Additional Information****Gene ID** 79065**Other Names**

Autophagy-related protein 9A, APG9-like 1, mATG9, ATG9A, APG9L1

**Target/Specificity**

Human ATG9A.

**Reconstitution & Storage**

Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles. Store undiluted.

**Precautions**

ATG9A Antibody (C-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

**ATG9A Antibody (C-Terminus) - Protein Information****Name** ATG9A {ECO:0000303|PubMed:20124090, ECO:0000312|HGNC:HGNC:22408}**Function**

Phospholipid scramblase involved in autophagy by mediating autophagosomal membrane expansion (PubMed:<a href="http://www.uniprot.org/citations/22456507" target="\_blank">22456507</a>, PubMed:<a href="http://www.uniprot.org/citations/27510922" target="\_blank">27510922</a>, PubMed:<a href="http://www.uniprot.org/citations/29437695" target="\_blank">29437695</a>, PubMed:<a href="http://www.uniprot.org/citations/32513819" target="\_blank">32513819</a>, PubMed:<a href="http://www.uniprot.org/citations/32610138" target="\_blank">32610138</a>, PubMed:<a href="http://www.uniprot.org/citations/33106659" target="\_blank">33106659</a>, PubMed:<a href="http://www.uniprot.org/citations/33468622" target="\_blank">33468622</a>, PubMed:<a href="http://www.uniprot.org/citations/33850023" target="\_blank">33850023</a>). Cycles between the preautophagosomal structure/phagophore

assembly site (PAS) and the cytoplasmic vesicle pool and supplies membrane for the growing autophagosome (PubMed:<a href="http://www.uniprot.org/citations/16940348" target="\_blank">16940348</a>, PubMed:<a href="http://www.uniprot.org/citations/22456507" target="\_blank">22456507</a>, PubMed:<a href="http://www.uniprot.org/citations/33106659" target="\_blank">33106659</a>). Lipid scramblase activity plays a key role in preautophagosomal structure/phagophore assembly by distributing the phospholipids that arrive through ATG2 (ATG2A or ATG2B) from the cytoplasmic to the luminal leaflet of the bilayer, thereby driving autophagosomal membrane expansion (PubMed:<a href="http://www.uniprot.org/citations/33106659" target="\_blank">33106659</a>). Also required to supply phosphatidylinositol 4- phosphate to the autophagosome initiation site by recruiting the phosphatidylinositol 4-kinase beta (PI4KB) in a process dependent on ARFIP2, but not ARFIP1 (PubMed:<a href="http://www.uniprot.org/citations/30917996" target="\_blank">30917996</a>). In addition to autophagy, also plays a role in necrotic cell death (By similarity).

### **Cellular Location**

Preatophagosomal structure membrane; Multi-pass membrane protein. Cytoplasmic vesicle, autophagosome membrane; Multi- pass membrane protein. Golgi apparatus, trans-Golgi network membrane; Multi-pass membrane protein. Late endosome membrane; Multi-pass membrane protein. Recycling endosome membrane; Multi-pass membrane protein. Endoplasmic reticulum membrane; Multi-pass membrane protein. Mitochondrion membrane; Multi-pass membrane protein. Note=Mainly localizes to the trans-Golgi network (TGN) and the endosomal system; cycles between them though vesicle trafficking (PubMed:27316455, PubMed:27663665). Export from the TGN to promote formation of autophagosomes is mediated by the AP-4 complex (PubMed:29180427, PubMed:30262884). Under amino acid starvation or rapamycin treatment, redistributes to preautophagosomal structure/phagophore assembly site (PAS) (PubMed:16940348). The starvation-induced redistribution depends on ULK1, ATG13, as well as SH3GLB1 (PubMed:16940348). Upon autophagy induction, a small portion transiently localizes to the autophagic membranes (PubMed:22456507) Recruited to damaged mitochondria during mitophagy in a RIMOC1- dependent manner (PubMed:34432599).

### **Volume**


50 µl

### **ATG9A Antibody (C-Terminus) - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


### **ATG9A Antibody (C-Terminus) - Images**



Immunofluorescence of ATG9A in human heart tissue with ATG9A antibody at 20 ug/ml.



Western blot analysis of ATG9A in mouse heart tissue lysate with ATG9A antibody at 1 ug/ml.



Anti-ATG9A antibody IHC of human spleen.

#### ATG9A Antibody (C-Terminus) - Background

Involved in autophagy and cytoplasm to vacuole transport (Cvt) vesicle formation. Plays a key role in the organization of the preautophagosomal structure/phagophore assembly site (PAS), the nucleating site for formation of the sequestering vesicle. Cycles between a juxta-nuclear trans-Golgi network compartment and late endosomes. Nutrient starvation induces accumulation on

autophagosomes. Starvation-dependent trafficking requires ULK1, ATG13 and Supt20H.

#### **ATG9A Antibody (C-Terminus) - References**

- Bechtel S.,et al.BMC Genomics 8:399-399(2007).  
Hillier L.W.,et al.Nature 434:724-731(2005).  
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.  
Ota T.,et al.Nat. Genet. 36:40-45(2004).  
Yamada T.,et al.J. Biol. Chem. 280:18283-18290(2005).