

SLC39A14 / ZIP14 Antibody (Internal)
Rabbit Polyclonal Antibody
Catalog # ALS14735**Specification****SLC39A14 / ZIP14 Antibody (Internal) - Product Information**

Application	WB, IHC-P, IF
Primary Accession	Q15043
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	54kDa KDa
Dilution	WB~~1:1000 IHC-P~~N/A IF~~1:50~200

SLC39A14 / ZIP14 Antibody (Internal) - Additional Information**Gene ID** 23516**Other Names**

Zinc transporter ZIP14, LIV-1 subfamily of ZIP zinc transporter 4, LZT-Hs4, Solute carrier family 39 member 14, Zrt- and Irt-like protein 14, ZIP-14, SLC39A14, KIAA0062, ZIP14

Target/Specificity

Human SLC39A14. At least three isoforms of ZIP14 are known to exist; this antibody will detect both isoforms. ZIP14 antibody is predicted to not cross-react with other ZIP family members.

Reconstitution & Storage

Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles. Store undiluted.

Precautions

SLC39A14 / ZIP14 Antibody (Internal) is for research use only and not for use in diagnostic or therapeutic procedures.

SLC39A14 / ZIP14 Antibody (Internal) - Protein Information**Name** SLC39A14 ([HGNC:20858](#))**Function**

Electroneutral transporter of the plasma membrane mediating the cellular uptake of the divalent metal cations zinc, manganese and iron that are important for tissue homeostasis, metabolism, development and immunity (PubMed:15642354, PubMed:27231142, PubMed:29621230). Functions as an energy-dependent symporter, transporting through the membranes an electroneutral complex composed of a divalent metal cation and two bicarbonate anions (By similarity). Beside these endogenous cellular substrates, can also import

cadmium a non-essential metal which is cytotoxic and carcinogenic (By similarity). Controls the cellular uptake by the intestinal epithelium of systemic zinc, which is in turn required to maintain tight junctions and the intestinal permeability (By similarity). Modifies the activity of zinc-dependent phosphodiesterases, thereby indirectly regulating G protein-coupled receptor signaling pathways important for gluconeogenesis and chondrocyte differentiation (By similarity). Regulates insulin receptor signaling, glucose uptake, glycogen synthesis and gluconeogenesis in hepatocytes through the zinc-dependent intracellular catabolism of insulin (PubMed:27703010). Through zinc cellular uptake also plays a role in the adaptation of cells to endoplasmic reticulum stress (By similarity). Major manganese transporter of the basolateral membrane of intestinal epithelial cells, it plays a central role in manganese systemic homeostasis through intestinal manganese uptake (PubMed:31028174). Also involved in manganese extracellular uptake by cells of the blood-brain barrier (PubMed:31699897). May also play a role in manganese and zinc homeostasis participating in their elimination from the blood through the hepatobiliary excretion (By similarity). Also functions in the extracellular uptake of free iron. May also function intracellularly and mediate the transport from endosomes to cytosol of iron endocytosed by transferrin (PubMed:20682781). Plays a role in innate immunity by regulating the expression of cytokines by activated macrophages (PubMed:23052185).

Cellular Location

Cell membrane; Multi-pass membrane protein. Apical cell membrane; Multi-pass membrane protein. Basolateral cell membrane; Multi-pass membrane protein. Early endosome membrane; Multi-pass membrane protein. Late endosome membrane; Multi-pass membrane protein. Lysosome membrane; Multi-pass membrane protein. Note=Localized and functional at both apical and basolateral membranes of microvascular capillary endothelial cells that constitute the blood-brain barrier (PubMed:31699897). Localized at the basolateral membrane of enterocytes (PubMed:31028174). Enriched at the plasma membrane upon glucose uptake (PubMed:27703010).

Tissue Location

Ubiquitously expressed, with higher expression in liver, pancreas, fetal liver, thyroid gland, left and right ventricle, right atrium and fetal heart (PubMed:15642354, PubMed:20682781, PubMed:7584044). Weakly expressed in spleen, thymus, and peripheral blood leukocytes (PubMed:7584044). Expressed in liver and in brain by large neurons in the globus pallidus, the insular cortex and the dentate nucleus and to a lower extent in the putamen and the caudate nucleus (at protein level) (PubMed:27231142). Expressed in osteoblasts and giant osteoclast-like cells, but not in osteocytes found osteoblastoma and giant cell tumors (at protein level) (PubMed:29621230). Expressed by microvascular capillary endothelial cells that constitute the blood-brain barrier (at protein level) (PubMed:31699897). Expressed by macrophages (PubMed:23052185)

Volume

50 µl


SLC39A14 / ZIP14 Antibody (Internal) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)

- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


SLC39A14 / ZIP14 Antibody (Internal) - Images



Immunofluorescence of ZIP14 in Mouse Liver cells with ZIP14 antibody at 20 ug/ml.



Anti-SLC39A14 antibody IHC of human colon, vessels.



Anti-SLC39A14 antibody IHC of human kidney.

SLC39A14 / ZIP14 Antibody (Internal) - Background

May mediate cellular uptake of nontransferrin-bound iron (By similarity). Broad-scope metal ion

transporter with a preference for zinc uptake.

SLC39A14 / ZIP14 Antibody (Internal) - References

Nomura N.,et al.DNA Res. 1:223-229(1994).

Ota T.,et al.Nat. Genet. 36:40-45(2004).

Nusbaum C.,et al.Nature 439:331-335(2006).

Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.

Taylor K.M.,et al.Biochim. Biophys. Acta 1611:16-30(2003).