

**FOS / c-FOS Antibody (aa1-50)**  
**Rabbit Polyclonal Antibody**  
**Catalog # ALS14745****Specification**

---

**FOS / c-FOS Antibody (aa1-50) - Product Information**

Application	WB, IHC-P, E
Primary Accession	<a href="#">P01100</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	41kDa KDa
Dilution	WB~~1:1000 IHC-P~~N/A E~~N/A

**FOS / c-FOS Antibody (aa1-50) - Additional Information****Gene ID** 2353**Other Names**

Proto-oncogene c-Fos, Cellular oncogene fos, G0/G1 switch regulatory protein 7, FOS, G0S7

**Target/Specificity**

Fos Antibody detects endogenous levels of total Fos protein.

**Reconstitution & Storage**

Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles.

**Precautions**

FOS / c-FOS Antibody (aa1-50) is for research use only and not for use in diagnostic or therapeutic procedures.

**FOS / c-FOS Antibody (aa1-50) - Protein Information****Name** FOS**Synonyms** G0S7**Function**

Nuclear phosphoprotein which forms a tight but non-covalently linked complex with the JUN/AP-1 transcription factor. In the heterodimer, FOS and JUN/AP-1 basic regions each seems to interact with symmetrical DNA half sites. On TGF-beta activation, forms a multimeric SMAD3/SMAD4/JUN/FOS complex at the AP1/SMAD-binding site to regulate TGF-beta-mediated signaling. Has a critical function in regulating the development of cells destined to form and maintain the skeleton. It is thought to have an important role in signal transduction, cell proliferation and differentiation. In growing cells, activates phospholipid synthesis, possibly by activating CDS1 and PI4K2A. This activity requires Tyr-dephosphorylation and association with the

endoplasmic reticulum.

#### Cellular Location

Nucleus. Endoplasmic reticulum. Cytoplasm, cytosol. Note=In quiescent cells, present in very small amounts in the cytosol. Following induction of cell growth, first localizes to the endoplasmic reticulum and only later to the nucleus. Localization at the endoplasmic reticulum requires dephosphorylation at Tyr-10 and Tyr- 30

#### Volume


Array

### FOS / c-FOS Antibody (aa1-50) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### FOS / c-FOS Antibody (aa1-50) - Images



Western blot of extracts from HepG2 cells, using Fos Antibody.

### FOS / c-FOS Antibody (aa1-50) - Background

Nuclear phosphoprotein which forms a tight but non-covalently linked complex with the JUN/AP-1 transcription factor. In the heterodimer, FOS and JUN/AP-1 basic regions each seems to interact with symmetrical DNA half sites. On TGF-beta activation, forms a multimeric SMAD3/SMAD4/JUN/FOS complex at the AP1/SMAD-binding site to regulate TGF-beta-mediated signaling. Has a critical function in regulating the development of cells destined to form and maintain the skeleton. It is thought to have an important role in signal transduction, cell proliferation and differentiation. In growing cells, activates phospholipid synthesis, possibly by activating CDS1 and PI4K2A. This activity requires Tyr-dephosphorylation and association with the endoplasmic reticulum.

### FOS / c-FOS Antibody (aa1-50) - References

van Straaten F., et al. Proc. Natl. Acad. Sci. U.S.A. 80:3183-3187(1983).  
Ota T., et al. Nat. Genet. 36:40-45(2004).  
Heilig R., et al. Nature 421:601-607(2003).

Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.  
Roux P.,et al.Oncogene 6:2155-2160(1991).