

**DNMT3A Antibody (aa170-220)**  
**Rabbit Polyclonal Antibody**  
**Catalog # ALS14907****Specification**

---

**DNMT3A Antibody (aa170-220) - Product Information**

Application	WB
Primary Accession	<a href="#">Q9Y6K1</a>
Reactivity	Human, Mouse, Chicken, Horse, Bovine, Cat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	102kDa KDa

**DNMT3A Antibody (aa170-220) - Additional Information****Gene ID** 1788**Other Names**

DNA (cytosine-5)-methyltransferase 3A, Dnmt3a, 2.1.1.37, DNA methyltransferase HsaIIIA, DNA MTase HsaIIIA, M.HsaIIIA, DNMT3A

**Target/Specificity**

Human DNMT3A

**Reconstitution & Storage**

Store at 4°C for short term applications. For long term storage, aliquot and store at -20°C.

**Precautions**

DNMT3A Antibody (aa170-220) is for research use only and not for use in diagnostic or therapeutic procedures.

**DNMT3A Antibody (aa170-220) - Protein Information****Name** DNMT3A**Function**

Required for genome-wide de novo methylation and is essential for the establishment of DNA methylation patterns during development (PubMed:<a href="http://www.uniprot.org/citations/12138111" target="\_blank">12138111</a>, PubMed:<a href="http://www.uniprot.org/citations/16357870" target="\_blank">16357870</a>, PubMed:<a href="http://www.uniprot.org/citations/30478443" target="\_blank">30478443</a>). DNA methylation is coordinated with methylation of histones (PubMed:<a href="http://www.uniprot.org/citations/12138111" target="\_blank">12138111</a>, PubMed:<a href="http://www.uniprot.org/citations/16357870" target="\_blank">16357870</a>, PubMed:<a href="http://www.uniprot.org/citations/30478443" target="\_blank">30478443</a>). It modifies DNA in a non-processive manner and also methylates non-CpG sites (PubMed:<a href="http://www.uniprot.org/citations/12138111" target="\_blank">12138111</a>, PubMed:<a href="http://www.uniprot.org/citations/16357870" target="\_blank">16357870</a>, PubMed:<a href="http://www.uniprot.org/citations/30478443" target="\_blank">30478443</a>).

href="http://www.uniprot.org/citations/16357870" target="\_blank">16357870</a>, PubMed:<a href="http://www.uniprot.org/citations/30478443" target="\_blank">30478443</a>). May preferentially methylate DNA linker between 2 nucleosomal cores and is inhibited by histone H1 (By similarity). Plays a role in paternal and maternal imprinting (By similarity). Required for methylation of most imprinted loci in germ cells (By similarity). Acts as a transcriptional corepressor for ZBTB18 (By similarity). Recruited to trimethylated 'Lys-36' of histone H3 (H3K36me3) sites (By similarity). Can actively repress transcription through the recruitment of HDAC activity (By similarity). Also has weak auto-methylation activity on Cys-710 in absence of DNA (By similarity).

#### Cellular Location

Nucleus. Chromosome Cytoplasm. Note=Accumulates in the major satellite repeats at pericentric heterochromatin {ECO:0000250|UniProtKB:O88508}

#### Tissue Location


Highly expressed in fetal tissues, skeletal muscle, heart, peripheral blood mononuclear cells, kidney, and at lower levels in placenta, brain, liver, colon, spleen, small intestine and lung

### DNMT3A Antibody (aa170-220) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### DNMT3A Antibody (aa170-220) - Images



Western blot of Dnmt3a in transfected 293 cell lysate, using ALS14907 at 1 ug/ml.

### DNMT3A Antibody (aa170-220) - Background

Required for genome-wide de novo methylation and is essential for the establishment of DNA methylation patterns during development. DNA methylation is coordinated with methylation of histones. It modifies DNA in a non-processive manner and also methylates non-CpG sites. May preferentially methylate DNA linker between 2 nucleosomal cores and is inhibited by histone H1.

Plays a role in paternal and maternal imprinting. Required for methylation of most imprinted loci in germ cells. Acts as a transcriptional corepressor for ZBTB18. Recruited to trimethylated 'Lys-36' of histone H3 (H3K36me3) sites. Can actively repress transcription through the recruitment of HDAC activity.

#### **DNMT3A Antibody (aa170-220) - References**

Xie S.,et al.Gene 236:87-95(1999).  
Chen T.,et al.J. Biol. Chem. 277:38746-38754(2002).  
Kim G.-D.,et al.EMBO J. 21:4183-4195(2002).  
Hillier L.W.,et al.Nature 434:724-731(2005).  
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.