

NCAM2 Antibody (C-Terminus)
Goat Polyclonal Antibody
Catalog # ALS14946**Specification**

NCAM2 Antibody (C-Terminus) - Product Information

Application	IHC
Primary Accession	O15394
Reactivity	Human, Rabbit, Hamster, Monkey, Horse, Bovine, Dog
Host	Goat
Clonality	Polyclonal
Calculated MW	93kDa KDa

NCAM2 Antibody (C-Terminus) - Additional Information**Gene ID** 4685**Other Names**

Neural cell adhesion molecule 2, N-CAM-2, NCAM-2, NCAM2, NCAM21

Target/Specificity

Human NCAM2.

Reconstitution & Storage

Store at -20°C. Minimize freezing and thawing.

Precautions

NCAM2 Antibody (C-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

NCAM2 Antibody (C-Terminus) - Protein Information**Name** NCAM2**Synonyms** NCAM21**Function**

May play important roles in selective fasciculation and zone- to-zone projection of the primary olfactory axons.

Cellular Location

Cell membrane; Single-pass type I membrane protein

Tissue Location

Expressed most strongly in adult and fetal brain.

Volume


50 µl

NCAM2 Antibody (C-Terminus) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NCAM2 Antibody (C-Terminus) - Images



Anti-NCAM2 antibody IHC of human brain, cortex.

NCAM2 Antibody (C-Terminus) - Background

May play important roles in selective fasciculation and zone-to-zone projection of the primary olfactory axons.

NCAM2 Antibody (C-Terminus) - References

Paoloni-Giacobino A., et al. Genomics 43:43-51(1997).
Ota T., et al. Nat. Genet. 36:40-45(2004).
Hattori M., et al. Nature 405:311-319(2000).
Zhang H., et al. Nat. Biotechnol. 21:660-666(2003).