

NPM1 / NPM / Nucleophosmin Antibody (aa201-250)
Rabbit Polyclonal Antibody
Catalog # ALS15004**Specification**

NPM1 / NPM / Nucleophosmin Antibody (aa201-250) - Product Information

Application	IF, WB, IHC
Primary Accession	P06748
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	33kDa KDa

NPM1 / NPM / Nucleophosmin Antibody (aa201-250) - Additional Information**Gene ID** 4869**Other Names**

Nucleophosmin, NPM, Nucleolar phosphoprotein B23, Nucleolar protein NO38, Numatrin, NPM1, NPM

Target/Specificity

NPM (Ab-234) Antibody detects endogenous levels of total NPM protein.

Reconstitution & Storage

Store at -20°C for up to one year.

Precautions

NPM1 / NPM / Nucleophosmin Antibody (aa201-250) is for research use only and not for use in diagnostic or therapeutic procedures.

NPM1 / NPM / Nucleophosmin Antibody (aa201-250) - Protein Information**Name** NPM1**Synonyms** NPM**Function**

Involved in diverse cellular processes such as ribosome biogenesis, centrosome duplication, protein chaperoning, histone assembly, cell proliferation, and regulation of tumor suppressors p53/TP53 and ARF. Binds ribosome presumably to drive ribosome nuclear export. Associated with nucleolar ribonucleoprotein structures and bind single-stranded nucleic acids. Acts as a chaperonin for the core histones H3, H2B and H4. Stimulates APEX1 endonuclease activity on apurinic/apyrimidinic (AP) double-stranded DNA but inhibits APEX1 endonuclease activity on AP single-stranded RNA. May exert a control of APEX1 endonuclease activity within nucleoli devoted to repair AP on rDNA and the removal of oxidized rRNA molecules. In concert with BRCA2, regulates centrosome duplication. Regulates centriole duplication: phosphorylation by PLK2 is able to trigger centriole replication. Negatively regulates the activation of EIF2AK2/PKR and suppresses

apoptosis through inhibition of EIF2AK2/PKR autophosphorylation. Antagonizes the inhibitory effect of ATF5 on cell proliferation and relieves ATF5-induced G2/M blockade (PubMed:22528486). In complex with MYC enhances the transcription of MYC target genes (PubMed:25956029). May act as chaperonin or cotransporter in the nucleolar localization of transcription termination factor TTF1 (By similarity).

Cellular Location

Nucleus, nucleolus. Nucleus, nucleoplasm. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome Note=Generally nucleolar, but is translocated to the nucleoplasm in case of serum starvation or treatment with anticancer drugs. Has been found in the cytoplasm in patients with primary acute myelogenous leukemia (AML), but not with secondary AML. Can shuttle between cytoplasm and nucleus. Co- localizes with the methylated form of RPS10 in the granular component (GC) region of the nucleolus. Colocalized with nucleolin and APEX1 in nucleoli. Isoform 1 of NEK2 is required for its localization to the centrosome during mitosis

Volume


50 µl

NPM1 / NPM / Nucleophosmin Antibody (aa201-250) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


NPM1 / NPM / Nucleophosmin Antibody (aa201-250) - Images



Immunofluorescence of HeLa cells, using NPM (Ab-234) Antibody.



Western blot of extracts from HeLa/NIH-3T3 cells, using NPM (Ab-234) Antibody.



Anti-NPM1 antibody IHC of human testis.

NPM1 / NPM / Nucleophosmin Antibody (aa201-250) - Background

Involved in diverse cellular processes such as ribosome biogenesis, centrosome duplication, protein chaperoning, histone assembly, cell proliferation, and regulation of tumor suppressors p53/TP53 and ARF. Binds ribosome presumably to drive ribosome nuclear export. Associated with nucleolar ribonucleoprotein structures and bind single-stranded nucleic acids. Acts as a chaperonin for the core histones H3, H2B and H4. Stimulates APEX1 endonuclease activity on apurinic/apyrimidinic (AP) double-stranded DNA but inhibits APEX1 endonuclease activity on AP single-stranded RNA. May exert a control of APEX1 endonuclease activity within nucleoli devoted to repair AP on rDNA and the removal of oxidized rRNA molecules. In concert with BRCA2, regulates centrosome duplication. Regulates centriole duplication: phosphorylation by PLK2 is able to trigger centriole replication. Negatively regulates the activation of EIF2AK2/PKR and suppresses apoptosis through inhibition of EIF2AK2/PKR autophosphorylation.

NPM1 / NPM / Nucleophosmin Antibody (aa201-250) - References

- Chan W.-Y., et al. *Biochemistry* 28:1033-1039(1989).
Li X., et al. *Biochem. Biophys. Res. Commun.* 163:72-78(1989).
Zhang X.T., et al. *Biochem. Biophys. Res. Commun.* 164:176-184(1989).
Chan P.-K., et al. *Nucleic Acids Res.* 25:1225-1232(1997).
Okuwaki M., et al. Submitted (APR-2000) to the EMBL/GenBank/DBJ databases.