

CFL1 / Cofilin Antibody (clone AF2C2)
Mouse Monoclonal Antibody
Catalog # ALS15197**Specification**

CFL1 / Cofilin Antibody (clone AF2C2) - Product Information

Application	WB, IHC
Primary Accession	P23528
Reactivity	Human, Mouse, Rat
Host	Mouse
Clonality	Monoclonal
Calculated MW	19kDa KDa

CFL1 / Cofilin Antibody (clone AF2C2) - Additional Information**Gene ID** 1072**Other Names**

Cofilin-1, 18 kDa phosphoprotein, p18, Cofilin, non-muscle isoform, CFL1, CFL

Reconstitution & Storage

Long term: -20°C; Short term: -20°C

Precautions

CFL1 / Cofilin Antibody (clone AF2C2) is for research use only and not for use in diagnostic or therapeutic procedures.

CFL1 / Cofilin Antibody (clone AF2C2) - Protein Information**Name** CFL1**Synonyms** CFL**Function**

Binds to F-actin and exhibits pH-sensitive F-actin depolymerizing activity (PubMed:11812157). In conjunction with the subcortical maternal complex (SCMC), plays an essential role for zygotes to progress beyond the first embryonic cell divisions via regulation of actin dynamics (PubMed:15580268). Required for the centralization of the mitotic spindle and symmetric division of zygotes (By similarity). Plays a role in the regulation of cell morphology and cytoskeletal organization in epithelial cells (PubMed:21834987). Required for the up-regulation of atypical chemokine receptor ACKR2 from endosomal compartment to cell membrane, increasing its efficiency in chemokine uptake and degradation (PubMed:23633677). Required for neural tube morphogenesis and neural crest cell migration (By similarity).

Cellular Location

Nucleus matrix. Cytoplasm, cytoskeleton. Cell projection, ruffle membrane; Peripheral membrane protein; Cytoplasmic side. Cell projection, lamellipodium membrane; Peripheral membrane protein; Cytoplasmic side. Cell projection, lamellipodium {ECO:0000250|UniProtKB:P18760}. Cell projection, growth cone {ECO:0000250|UniProtKB:P18760}. Cell projection, axon {ECO:0000250|UniProtKB:P18760}. Note=Colocalizes with the actin cytoskeleton in membrane ruffles and lamellipodia. Detected at the cleavage furrow and contractile ring during cytokinesis. Almost completely in nucleus in cells exposed to heat shock or 10% dimethyl sulfoxide

Tissue Location

Widely distributed in various tissues.

Volume


50 µl

CFL1 / Cofilin Antibody (clone AF2C2) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CFL1 / Cofilin Antibody (clone AF2C2) - Images



0



Anti-Cofilin antibody IHC of human thymus.

CFL1 / Cofilin Antibody (clone AF2C2) - Background

Binds to F-actin and exhibits pH-sensitive F-actin depolymerizing activity. Regulates actin cytoskeleton dynamics. Important for normal progress through mitosis and normal cytokinesis. Plays a role in the regulation of cell morphology and cytoskeletal organization. Required for the up-regulation of atypical chemokine receptor ACKR2 from endosomal compartment to cell membrane, increasing its efficiency in chemokine uptake and degradation.

CFL1 / Cofilin Antibody (clone AF2C2) - References

Ogawa K., et al. Nucleic Acids Res. 18:7169-7169(1990).
van der Steege G., et al. Eur. J. Hum. Genet. 3:87-95(1995).
Gillett G.T., et al. Ann. Hum. Genet. 60:201-211(1996).
Kalnine N., et al. Submitted (MAY-2003) to the EMBL/GenBank/DDBJ databases.
Ota T., et al. Nat. Genet. 36:40-45(2004).