

GRIK1 / GLUR5 Antibody (Internal)

Goat Polyclonal Antibody Catalog # ALS15357

Specification

GRIK1 / GLUR5 Antibody (Internal) - Product Information

Application WB, IHC Primary Accession P39086

Reactivity Human, Mouse, Rat, Monkey, Horse

Host Goat
Clonality Polyclonal
Calculated MW 104kDa KDa

GRIK1 / GLUR5 Antibody (Internal) - Additional Information

Gene ID 2897

Other Names

Glutamate receptor ionotropic, kainate 1, GluK1, Excitatory amino acid receptor 3, EAA3, Glutamate receptor 5, GluR-5, GluR5, GRIK1, GLUR5

Target/Specificity

Human GRIK1. This antibody is expected to recognise both reported isoforms according to NP 000821.1 and NP 783300.1.

Reconstitution & Storage

Store at -20°C. Minimize freezing and thawing.

Precautions

GRIK1 / GLUR5 Antibody (Internal) is for research use only and not for use in diagnostic or therapeutic procedures.

GRIK1 / GLUR5 Antibody (Internal) - Protein Information

Name GRIK1

Synonyms GLUR5

Function

Ionotropic glutamate receptor. L-glutamate acts as an excitatory neurotransmitter at many synapses in the central nervous system. Binding of the excitatory neurotransmitter L-glutamate induces a conformation change, leading to the opening of the cation channel, and thereby converts the chemical signal to an electrical impulse. The receptor then desensitizes rapidly and enters a transient inactive state, characterized by the presence of bound agonist. May be involved in the transmission of light information from the retina to the hypothalamus.

Cellular Location

Cell membrane; Multi-pass membrane protein. Postsynaptic cell membrane; Multi-pass membrane



protein

Tissue Location

Most abundant in the cerebellum and the suprachiasmatic nuclei (SCN) of the hypothalamus

GRIK1 / GLUR5 Antibody (Internal) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

GRIK1 / GLUR5 Antibody (Internal) - Images



Antibody (1 ug/ml) staining of Human Brain (Cerebellum) lysate (35 ug protein in RIPA buffer).



Anti-GRIK1 / GLUR5 antibody IHC of human brain, cortex.

GRIK1 / GLUR5 Antibody (Internal) - Background

lonotropic glutamate receptor. L-glutamate acts as an excitatory neurotransmitter at many synapses in the central nervous system. Binding of the excitatory neurotransmitter L- glutamate





induces a conformation change, leading to the opening of the cation channel, and thereby converts the chemical signal to an electrical impulse. The receptor then desensitizes rapidly and enters a transient inactive state, characterized by the presence of bound agonist. May be involved in the transmission of light information from the retina to the hypothalamus.

GRIK1 / GLUR5 Antibody (Internal) - References

Gregor P.,et al.NeuroReport 4:1343-1346(1993). Korczak B.,et al.Recept. Channels 3:41-49(1995). Langer A.,et al.Submitted (SEP-1999) to the EMBL/GenBank/DDBJ databases. Nutt S.L.,et al.NeuroReport 5:2625-2629(1994). Shibata H.,et al.Psychiatr. Genet. 11:139-144(2001).