

CYP2E1 Antibody (Internal)
Goat Polyclonal Antibody
Catalog # ALS15387**Specification**

CYP2E1 Antibody (Internal) - Product Information

Application	WB, IHC
Primary Accession	P05181
Reactivity	Human
Host	Goat
Clonality	Polyclonal
Calculated MW	57kDa KDa

CYP2E1 Antibody (Internal) - Additional Information**Gene ID** 1571**Other Names**

Cytochrome P450 2E1, 1.14.13.-, 4-nitrophenol 2-hydroxylase, 1.14.13.n7, CYP11E1, Cytochrome P450-J, Cytochrome P450 2E1, N-terminally processed, CYP2E1, CYP2E

Target/Specificity

Human CYP2E1.

Reconstitution & Storage

Store at -20°C. Minimize freezing and thawing.

Precautions

CYP2E1 Antibody (Internal) is for research use only and not for use in diagnostic or therapeutic procedures.

CYP2E1 Antibody (Internal) - Protein Information**Name** CYP2E1 {ECO:0000303|PubMed:10553002, ECO:0000312|HGNC:HGNC:2631}**Function**

A cytochrome P450 monooxygenase involved in the metabolism of fatty acids (PubMed:10553002, PubMed:18577768). Mechanistically, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons provided by NADPH via cytochrome P450 reductase (NADPH--hemoprotein reductase) (PubMed:10553002, PubMed:18577768). Catalyzes the hydroxylation of carbon-hydrogen bonds. Hydroxylates fatty acids specifically at the omega-1 position displaying the highest catalytic activity for saturated fatty acids (PubMed:10553002, PubMed:18577768). May be

involved in the oxidative metabolism of xenobiotics (Probable).

Cellular Location

Endoplasmic reticulum membrane {ECO:0000250|UniProtKB:P05182}; Peripheral membrane protein {ECO:0000250|UniProtKB:P05182}. Microsome membrane {ECO:0000250|UniProtKB:P05182}; Peripheral membrane protein {ECO:0000250|UniProtKB:P05182}. Mitochondrion inner membrane {ECO:0000250|UniProtKB:P05182}; Peripheral membrane protein {ECO:0000250|UniProtKB:P05182}. Note=Post-translationally targeted to mitochondria. TOMM70 is required for the translocation across the mitochondrial outer membrane. After translocation into the matrix, associates with the inner membrane as a membrane extrinsic protein {ECO:0000250|UniProtKB:P05182}

CYP2E1 Antibody (Internal) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CYP2E1 Antibody (Internal) - Images



CYP2E1 antibody (0.05 ug/ml) staining of Human Liver lysate (35 ug protein/ml in RIPA buffer).



Anti-CYP2E1 antibody IHC of human liver.

CYP2E1 Antibody (Internal) - Background

Metabolizes several precarcinogens, drugs, and solvents to reactive metabolites. Inactivates a number of drugs and xenobiotics and also bioactivates many xenobiotic substrates to their hepatotoxic or carcinogenic forms.

CYP2E1 Antibody (Internal) - References

- Song B.-J., et al. J. Biol. Chem. 261:16689-16697(1986).
Umeno M., et al. Biochemistry 27:9006-9013(1988).
Zhuge J., et al. Submitted (SEP-1999) to the EMBL/GenBank/DDBJ databases.
Deloukas P., et al. Nature 429:375-381(2004).
Mural R.J., et al. Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.