

NEK7 Antibody (Internal)
Goat Polyclonal Antibody
Catalog # ALS15390**Specification**

NEK7 Antibody (Internal) - Product Information

Application	WB, IHC-P, E
Primary Accession	Q8TDX7
Reactivity	Human, Mouse, Rat, Rabbit, Hamster, Monkey, Pig, Chicken, Horse, Bovine, Dog
Host	Goat
Clonality	Polyclonal
Calculated MW	35kDa KDa
Dilution	WB~~1:1000 IHC-P~~N/A E~~N/A

NEK7 Antibody (Internal) - Additional Information**Gene ID** 140609**Other Names**

Serine/threonine-protein kinase Nek7, 2.7.11.1, Never in mitosis A-related kinase 7, NimA-related protein kinase 7, NEK7

Target/Specificity

Human NEK7.

Reconstitution & Storage

Store at -20°C. Minimize freezing and thawing.

Precautions

NEK7 Antibody (Internal) is for research use only and not for use in diagnostic or therapeutic procedures.

NEK7 Antibody (Internal) - Protein Information**Name** NEK7 {ECO:0000303|PubMed:11701951, ECO:0000312|HGNC:HGNC:13386}**Function**

Protein kinase which plays an important role in mitotic cell cycle progression (PubMed:17101132, PubMed:19941817, PubMed:31409757). Required for microtubule nucleation activity of the centrosome, robust mitotic spindle formation and cytokinesis (PubMed:17586473, PubMed:19414596, PubMed:19941817).

target="_blank">19941817, PubMed:26522158, PubMed:31409757). Phosphorylates EML4 at 'Ser-146', promoting its dissociation from microtubules during mitosis which is required for efficient chromosome congression (PubMed:31409757). Phosphorylates RPS6KB1 (By similarity). Acts as an essential activator of the NLRP3 inflammasome assembly independently of its kinase activity (PubMed:26642356, PubMed:36442502, PubMed:39173637). Acts by unlocking NLRP3 following NLRP3 translocation into the microtubule organizing center (MTOC), relieving NLRP3 autoinhibition and promoting formation of the NLRP3:PYCARD complex, and activation of CASP1 (PubMed:26642356, PubMed:31189953, PubMed:36442502, PubMed:39173637). Serves as a cellular switch that enforces mutual exclusivity of the inflammasome response and cell division: interaction with NEK9 prevents interaction with NLRP3 and activation of the inflammasome during mitosis (PubMed:26642356, PubMed:31189953).

Cellular Location

Nucleus {ECO:0000250|UniProtKB:Q9ES74}. Cytoplasm. Cytoplasm, cytoskeleton, spindle pole. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Note=Present at centrosome throughout the cell cycle (PubMed:17586473). Also detected at spindle midzone of the anaphase cells and eventually concentrates at the midbody (PubMed:17586473). Interaction with ANKS3 prevents its translocation to the nucleus (By similarity). {ECO:0000250|UniProtKB:Q9ES74, ECO:0000269|PubMed:17586473}

Tissue Location


Highly expressed in lung, muscle, testis, brain, heart, liver, leukocyte and spleen. Lower expression in ovary, prostate and kidney. No expression seen in small intestine


NEK7 Antibody (Internal) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NEK7 Antibody (Internal) - Images





NEK7 antibody (0.05 ug/ml) staining of HeLa lysate (35 ug protein/ml in RIPA buffer).

NEK7 Antibody (Internal) - Background

Protein kinase which plays an important role in mitotic cell cycle progression. Required for microtubule nucleation activity of the centrosome, robust mitotic spindle formation and cytokinesis. Phosphorylates RPS6KB1.

NEK7 Antibody (Internal) - References

Kimura M.,et al.Cytogenet. Cell Genet. 94:33-38(2001).
Melton D.,et al.Submitted (MAR-2004) to the EMBL/GenBank/DDBJ databases.
Gregory S.G.,et al.Nature 441:315-321(2006).
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.
Belham C.,et al.J. Biol. Chem. 278:34897-34909(2003).