

**NEK7 Antibody (Internal)**  
**Goat Polyclonal Antibody**  
**Catalog # ALS15390****Specification****NEK7 Antibody (Internal) - Product Information**

Application	WB, IHC-P, E
Primary Accession	<a href="#">Q8TDX7</a>
Reactivity	Human, Mouse, Rat, Rabbit, Hamster, Monkey, Pig, Chicken, Horse, Bovine, Dog
Host	Goat
Clonality	Polyclonal
Calculated MW	35kDa KDa
Dilution	WB~~1:1000 IHC-P~~N/A E~~N/A

**NEK7 Antibody (Internal) - Additional Information****Gene ID** 140609**Other Names**

Serine/threonine-protein kinase Nek7, 2.7.11.1, Never in mitosis A-related kinase 7, NimA-related protein kinase 7, NEK7

**Target/Specificity**

Human NEK7.

**Reconstitution & Storage**

Store at -20°C. Minimize freezing and thawing.

**Precautions**

NEK7 Antibody (Internal) is for research use only and not for use in diagnostic or therapeutic procedures.

**NEK7 Antibody (Internal) - Protein Information****Name** NEK7 {ECO:0000303|PubMed:11701951, ECO:0000312|HGNC:HGNC:13386}**Function**

Protein kinase which plays an important role in mitotic cell cycle progression (PubMed:<a href="http://www.uniprot.org/citations/17101132" target="\_blank">17101132</a>, PubMed:<a href="http://www.uniprot.org/citations/19941817" target="\_blank">19941817</a>, PubMed:<a href="http://www.uniprot.org/citations/31409757" target="\_blank">31409757</a>). Required for microtubule nucleation activity of the centrosome, robust mitotic spindle formation and cytokinesis (PubMed:<a href="http://www.uniprot.org/citations/17586473" target="\_blank">17586473</a>, PubMed:<a href="http://www.uniprot.org/citations/19414596" target="\_blank">19414596</a>, PubMed:<a href="http://www.uniprot.org/citations/19941817" target="\_blank">19941817</a>).

target="\_blank">>19941817</a>, PubMed:<a href="http://www.uniprot.org/citations/26522158" target="\_blank">26522158</a>, PubMed:<a href="http://www.uniprot.org/citations/31409757" target="\_blank">31409757</a>). Phosphorylates EML4 at 'Ser-146', promoting its dissociation from microtubules during mitosis which is required for efficient chromosome congression (PubMed:<a href="http://www.uniprot.org/citations/31409757" target="\_blank">31409757</a>). Phosphorylates RPS6KB1 (By similarity). Acts as an essential activator of the NLRP3 inflammasome assembly independently of its kinase activity (PubMed:<a href="http://www.uniprot.org/citations/26642356" target="\_blank">26642356</a>, PubMed:<a href="http://www.uniprot.org/citations/36442502" target="\_blank">36442502</a>, PubMed:<a href="http://www.uniprot.org/citations/39173637" target="\_blank">39173637</a>). Acts by unlocking NLRP3 following NLRP3 translocation into the microtubule organizing center (MTOC), relieving NLRP3 autoinhibition and promoting formation of the NLRP3:PYCARD complex, and activation of CASP1 (PubMed:<a href="http://www.uniprot.org/citations/26642356" target="\_blank">26642356</a>, PubMed:<a href="http://www.uniprot.org/citations/31189953" target="\_blank">31189953</a>, PubMed:<a href="http://www.uniprot.org/citations/36442502" target="\_blank">36442502</a>, PubMed:<a href="http://www.uniprot.org/citations/39173637" target="\_blank">39173637</a>). Serves as a cellular switch that enforces mutual exclusivity of the inflammasome response and cell division: interaction with NEK9 prevents interaction with NLRP3 and activation of the inflammasome during mitosis (PubMed:<a href="http://www.uniprot.org/citations/26642356" target="\_blank">26642356</a>, PubMed:<a href="http://www.uniprot.org/citations/31189953" target="\_blank">31189953</a>).

### Cellular Location

Nucleus {ECO:0000250|UniProtKB:Q9ES74}. Cytoplasm. Cytoplasm, cytoskeleton, spindle pole. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Note=Present at centrosome throughout the cell cycle (PubMed:17586473). Also detected at spindle midzone of the anaphase cells and eventually concentrates at the midbody (PubMed:17586473). Interaction with ANKS3 prevents its translocation to the nucleus (By similarity). {ECO:0000250|UniProtKB:Q9ES74, ECO:0000269|PubMed:17586473}

### Tissue Location


Highly expressed in lung, muscle, testis, brain, heart, liver, leukocyte and spleen. Lower expression in ovary, prostate and kidney. No expression seen in small intestine

### NEK7 Antibody (Internal) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### NEK7 Antibody (Internal) - Images



NEK7 antibody (0.05 ug/ml) staining of HeLa lysate (35 ug protein/ml in RIPA buffer).

#### **NEK7 Antibody (Internal) - Background**

Protein kinase which plays an important role in mitotic cell cycle progression. Required for microtubule nucleation activity of the centrosome, robust mitotic spindle formation and cytokinesis. Phosphorylates RPS6KB1.

#### **NEK7 Antibody (Internal) - References**

- Kimura M., et al. Cytogenet. Cell Genet. 94:33-38(2001).  
Melton D., et al. Submitted (MAR-2004) to the EMBL/GenBank/DDBJ databases.  
Gregory S.G., et al. Nature 441:315-321(2006).  
Mural R.J., et al. Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.  
Belham C., et al. J. Biol. Chem. 278:34897-34909(2003).