

PLD4 / Phospholipase D4 Antibody (aa457-506)
Rabbit Polyclonal Antibody
Catalog # ALS15743**Specification****PLD4 / Phospholipase D4 Antibody (aa457-506) - Product Information**

Application	WB, IHC-P, E
Primary Accession	Q96BZ4
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	56kDa KDa
Dilution	WB~~1:1000 IHC-P~~N/A E~~N/A

PLD4 / Phospholipase D4 Antibody (aa457-506) - Additional Information**Gene ID** 122618**Other Names**

Phospholipase D4, PLD 4, 3.1.4.4, Choline phosphatase 4, Phosphatidylcholine-hydrolyzing phospholipase D4, PLD4, C14orf175

Target/Specificity

PLD4 Antibody detects endogenous levels of total PLD4 protein.

Reconstitution & Storage

Store at -20°C for up to one year.

Precautions

PLD4 / Phospholipase D4 Antibody (aa457-506) is for research use only and not for use in diagnostic or therapeutic procedures.

PLD4 / Phospholipase D4 Antibody (aa457-506) - Protein Information**Name** PLD4 {ECO:0000303|PubMed:30111894, ECO:0000312|HGNC:HGNC:23792}**Function**

5'→3' exonuclease that hydrolyzes the phosphodiester bond of single-stranded DNA (ssDNA) and RNA molecules to form nucleoside 3'-monophosphates and 5'-end 5'-hydroxy deoxyribonucleotide/ribonucleotide fragments (PubMed:30111894, PubMed:34620855, PubMed:38537643, PubMed:39423811). Partially redundant with PLD3, can cleave all four nucleotides displaying higher efficiency for ssDNA and RNA fragments initiated with uridine and guanosine residues and lower efficiency for

cytidine-initiated substrates (PubMed:30111894, PubMed:34620855, PubMed:38537643, PubMed:39423811). As a result, it does not always degrade polynucleotides to the single nucleotide level, it can stall at specific sites sparing certain fragments from exonucleolytic degradation (PubMed:30111894, PubMed:34620855, PubMed:38537643, PubMed:39423811). Processes self and pathogenic ssDNA and RNA molecules that reach the endolysosomal compartment via phagocytosis or autophagy and may serve as 'danger' signals for recognition by innate immune receptors such as toll-like receptors (TLRs) (PubMed:38697119). Degrades mitochondrial CpG-rich ssDNA fragments to prevent TLR9 activation and autoinflammatory response, but it can cleave viral RNA to generate ligands for TLR7 activation and initiate antiviral immune responses (PubMed:38697119). In plasmacytoid dendritic cells, it cooperates with endonuclease RNASET2 to release 2',3'-cyclic guanosine monophosphate (2',3'-cGMP), a potent stimulatory ligand for TLR7 (PubMed:38697119). Produces 2',3'-cGMPs and cytidine-rich RNA fragments that occupy TLR7 ligand-binding pockets and trigger a signaling- competent state (PubMed:38697119). Can exert polynucleotide phosphatase activity toward 5'-phosphorylated ssDNA substrates although at a slow rate (PubMed:38537643). Transphosphatidylase that catalyzes the exchange with R to S stereo-inversion of the glycerol moiety between (S,R)-lysophosphatidylglycerol (LPG) and monoacylglycerol (MAG) substrates to yield (S,S)-bis(monoacylglycero)phosphate (BMP) (PubMed:39423811). Can synthesize a variety of (S,S)-BMPs representing the main phospholipid constituent of lysosomal intraluminal vesicle (ILV) membranes that bind acid hydrolases for lipid degradation (PubMed:39423811). Regulates the homeostasis and interorganellar communication of the endolysosomal system with an overall impact on cellular removal of dysfunctional organelles via autophagy as well as proper protein and lipid turnover. May play a role in myotube formation in response to ER stress (By similarity).

Cellular Location

Endoplasmic reticulum membrane {ECO:0000250|UniProtKB:Q8BG07}; Single-pass type II membrane protein {ECO:0000250|UniProtKB:Q8BG07}. Golgi apparatus, trans-Golgi network membrane {ECO:0000250|UniProtKB:Q8BG07}; Single-pass type II membrane protein {ECO:0000250|UniProtKB:Q8BG07}. Nucleus {ECO:0000250|UniProtKB:Q8BG07}. Early endosome {ECO:0000250|UniProtKB:Q8BG07}. Cytoplasmic vesicle, phagosome {ECO:0000250|UniProtKB:Q8BG07}. Lysosome Note=Activation of microglia induces translocation of PLD4 from the nucleus to the phagosomes. {ECO:0000250|UniProtKB:Q8BG07}

Tissue Location

Expressed in plasmacytoid dendritic cells and monocytes (at protein level).

Volume


50 µl

PLD4 / Phospholipase D4 Antibody (aa457-506) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PLD4 / Phospholipase D4 Antibody (aa457-506) - Images



Anti-PLD4 / Phospholipase D4 antibody IHC staining of human kidney.



Western blot of extracts from Jurkat cells, using PLD4 Antibody.

PLD4 / Phospholipase D4 Antibody (aa457-506) - References

Clark H.F., et al. Genome Res. 13:2265-2270(2003).